**DIMACS Technical Report 97-11**
**March 1997**

# On the density of subgraphs in a graph with bounded independence number[1]

by

Pavel Valtr[2]

DIMACS Center, Rutgers University
P.O.Box 1179, Piscataway, NJ 08855, U.S.A.
e-mail: valtr@dimacs.rutgers.edu

Department of Applied Mathematics, Charles University
Malostranské nám. 25, 118 00 Praha 1, Czech Republic
e-mail: valtr@kam.ms.mff.cuni.cz

[1]Research was supported by Czech Republic Grant GAČR 0194/1996 and by Charles University grant No. 194/1996.
[2]Postdoctoral Fellow

---

DIMACS is a cooperative project of Rutgers University, Princeton University, AT&T Bell Laboratories and Bellcore.

DIMACS is an NSF Science and Technology Center, funded under contract STC–91–19999; and also receives support from the New Jersey Commission on Science and Technology.

**ABSTRACT**

Let $\sigma(n, m, k)$ be the largest number $\sigma \in [0, 1]$ such that any graph on $n$ vertices with independence number at most $m$ has a subgraph on $k$ vertices with at least $\sigma \cdot \binom{k}{2}$ edges. Up to a constant multiplicative factor, we determine $\sigma(n, m, k)$ for all $n, m, k$. For $\log n \leq m = k \leq n$, our result gives $\sigma(n, m, m) = \Theta(\frac{\log(n/m)}{m})$, which was conjectured by Alon.

# 1 Introduction

All logarithms in this paper are assumed to be in base $e$. Let $\alpha(G)$ denote the independence number of a graph $G$. Erdős [Er79] conjectured that every graph $G$ on $n$ vertices with $\alpha(G) < \lfloor\sqrt{n}\rfloor$ contains a subgraph on $\lfloor\sqrt{n}\rfloor$ vertices with at least $c \cdot \sqrt{n}\log n$ edges, where $c > 0$ is a constant independent of $n$. This conjecture was proved by Alon [Al96] who further conjectured [GS95] that, for $\log n \leq m \leq n$, every graph $G$ on $n$ vertices with $\alpha(G) < m$ contains a subgraph on $m$ vertices with at least $c \cdot m\log(n/m)$ edges. Alon [Al96] proved his conjecture for $\log n \leq m \leq C\log n$ and for $n^{\varepsilon} \leq m \leq n$, where $C, \varepsilon > 0$ are any two positive constants. Alon [Al96] also proved that if his conjecture is true, then the bound $c \cdot m\log(n/m)$ in it is optimal up to a constant factor. In this paper, we prove Alon's conjecture by showing the following more general results.

**Theorem 1** *Let $2 \leq k \leq n$ and $2 \leq m \leq n/2$. Let $\sigma(n,m,k)$ be the largest number $\sigma \in [0,1]$ such that any graph on $n$ vertices with $\alpha(G) \leq m$ has a subgraph on $k$ vertices with at least $\sigma \cdot \binom{k}{2}$ edges. Then*

$$\sigma(n,m,k) = \begin{cases} \Theta\left(\min\left\{\frac{\log(en/k)}{m}, 1\right\}\right), & \text{for } k \geq m, \\ \Theta\left(\min\left\{\frac{\log(en/k)}{k\log(em/k)}, 1\right\}\right), & \text{for } m \geq k. \end{cases}$$

*($\sigma(n,m,k) = \Theta(f(n,m,k))$ means that there are two positive constants $c_1, c_2 > 0$ such that $c_1 f(n,m,k) \leq \sigma(n,m,k) \leq c_2 f(n,m,k)$ for all feasible $n, m, k$.)*

**Observation 2** *Let $2 \leq k \leq n$ and $n/2 \leq m \leq n$. Let $\sigma(n,m,k)$ be defined as in Theorem 1. Then*

$$\sigma(n,m,k) = \frac{\min\{n-m, \lfloor k/2\rfloor\}}{\binom{k}{2}}.$$

Theorem 1 and Observation 2 quantitatively express the fact that any graph with no dense subgraphs has a large independence number. Alon's conjecture indeed follows from them, since for $m = k$ we get $\sigma(n,m,m) = \Theta\left(\min\{\frac{\log(en/m)}{m}, 1\}\right) = \Theta\left(\frac{\log(n/m)}{m}\right)$ (if $\log n \leq m \leq n/2$) or $\sigma(n,m,m) = \frac{\min\{n-m,\lfloor m/2\rfloor\}}{\binom{m}{2}} = \Theta\left(\frac{n-m}{m^2}\right) = \Theta\left(\frac{\log(n/m)}{m}\right)$ (if $n/2 \leq m \leq n$).

Both the lower and the upper bounds on $\sigma(n,m,k)$ are proved differently for $m \leq k$ than for $k \leq m$. In both cases ($m \leq k$ and $k \leq m$), the lower bound is proved by "algorithmic" methods and the upper bound by probabilistic methods (using Alon's proof in the special case $m = k$).

Our result is closely related to Ramsey theory. Recall that the classical (off-diagonal) Ramsey number $r(m,k)$ is defined as the smallest number $r$ such that, for any 2-coloring of edges of $K_r$ by red and blue, there is a subgraph (isomorphic to) $K_m$ all of whose edges are red or a subgraph $K_k$ all of whose edges are blue. One can ask how the number $r(m,k)$ changes if in the latter case one requires only some fraction of edges in $K_k$ to be blue. More precisely, let $n_\sigma(m,k)$ be the smallest number $n$ such that, for any 2-coloring of edges of $K_n$

by red and blue, there is a subgraph $K_m$ all of whose edges are red or a subgraph $K_k$ with at least $\sigma\binom{k}{2}$ blue edges. For $\sigma \in [0,1]$, Theorem 1 and Observation 2 show that

$$n_\sigma(m,k) \approx \begin{cases} k \cdot 2^{c\cdot\sigma m}, & \text{for } k \geq m, \\ k \cdot 2^{c\cdot\sigma k \log(em/k)}, & \text{for } m \geq k. \end{cases}$$

We plan to study this question in a more general setting in a forthcoming paper.

Our proofs give relatively good constants of proportionality in Theorem 1. However, for the sake of simplicity, we do not try to optimize them.

Theorem 1 and Observation 2 might be also stated as results about densities of graphs. The *density* of a graph $G = (V, E)$ is the number $\frac{|E|}{\binom{|V|}{2}}$, i.e. the number of edges in $G$ divided by the number of edges in the complete graph on the same vertex set. The graph density satisfies the following useful observation.

**Observation 3** *Let $G$ be a graph on $m$ vertices of density $\sigma$. Then, for any $2 \leq k \leq m$, there is an (induced) subgraph of $G$ on $k$ vertices of density at least $\sigma$.*

*Proof.* Standard averaging argument. □

Theorem 1 is proved in Sections 2 (the upper bound) and 3 (the lower bound). Observation 2 has a simple proof:

*Proof of Observation 2.* Let $m \geq n/2$. The upper bound

$$\sigma(n,m,k) \leq \frac{\min\{n-m, \lfloor k/2 \rfloor\}}{\binom{k}{2}}$$

can be obtained from a matching of size $n-m$ on $n$ vertices. To see the lower bound

$$\sigma(n,m,k) \geq \frac{\min\{n-m, \lfloor k/2 \rfloor\}}{\binom{k}{2}},$$

observe that any graph $G$ on $n$ vertices with $\alpha(G) \leq m$ has at least $n-m$ edges. Any $\min\{n-m, \lfloor k/2 \rfloor\}$ of them have at most $k$ vertices altogether. The lower bound follows. □

# 2 The upper bound

## 2.1 The case $m \geq k$

The random graph $G_{n,p}$ is defined as a random graph on $n$ labeled vertices obtained by picking each pair of vertices as an edge, randomly and independently, with probability $p$. Set

$$p = \frac{4}{m-1}\log(en/m).$$

As shown in [Al96, Prop. 3.1] by a "brute-force" argument, if $p \leq 1$ then the random graph $G_{n,p}$ satisfies $\alpha(G_{n,p}) < m$ with probability bigger than $\frac{1}{2}$. Consequently, the following lemma is the core of the proof of the upper bound in Theorem 1 for $m \geq k$:

**Lemma 4** *For any $n \geq m \geq k \geq 2$ such that $p = \frac{4}{m-1}\log(en/m)$ is not bigger than 1, the random graph $G_{n,p}$ has a subgraph on $k$ vertices of density at least $8e^2\frac{\log(en/k)}{k\log(em/k)}$ with probability smaller than $\frac{1}{2}$.*

Lemma 4 gives the upper bound in Theorem 1 for $m \geq k$ and $p = \frac{4}{m-1}\log(en/m) \leq 1$ (indeed, in this case Lemma 4 and Alon's result mentioned above it show that the graph $G_{n,p}$ demonstrates the bound with positive probability). If $m \geq k$ and $p > 1$, then $m < 4\log(en/m) + 1 \leq 5\log(en/k)$, and consequently $\frac{\log(en/k)}{k\log(em/k)} \geq \frac{\log(en/k)}{m} > \frac{1}{5}$. Thus, for $m \geq k$ and $p > 1$ Theorem 1 asserts that $\sigma(n, m, k) = \Theta(1)$, and the upper bound in it is trivial in this case. It remains to prove Lemma 4.

In the proof of Lemma 4, we use the following estimate on binomial coefficients.

**Lemma 5** *If $n \geq k \geq 1$, then*

$$\binom{n}{k} < \left(\frac{en}{k}\right)^k.$$

Lemma 5 appears relatively often in the literature. We include its short proof for the sake of completeness.

*Proof of Lemma 5.* We fix $n$ and proceed by induction on $k$. For $k = 1$ the lemma trivially holds. Suppose now that $k > 1$ and that $\binom{n}{k-1} < \left(\frac{en}{k-1}\right)^{k-1}$. Then

$$\binom{n}{k} = \frac{n-k+1}{k}\binom{n}{k-1} < \frac{n}{k}\left(\frac{en}{k-1}\right)^{k-1} = \frac{1}{e}\left(\frac{k}{k-1}\right)^{k-1}\left(\frac{en}{k}\right)^k < \left(\frac{en}{k}\right)^k.$$

□

*Proof of Lemma 4.* Set $z = \left\lceil \frac{8e^2 \cdot \log(en/k)}{k\log(em/k)}\binom{k}{2} \right\rceil$. We will show (also by a "brute-force" argument) that the graph $G = G_{n,p}$ has a subgraph on $k$ vertices with at least $z$ edges with probability smaller than $\frac{1}{2}$.

There are $\binom{n}{k}$ sets of $k$ vertices of $G$. For any set $V_0$ of $k$ vertices of $G$, there are $\binom{\binom{k}{2}}{z}$ collections (sets) of $z$ pairs of vertices of $V_0$. Thus, there are at most $\binom{n}{k}\binom{\binom{k}{2}}{z}$ collections of $z$ pairs of vertices chosen from among at most $k$ distinct vertices in $G$. Denote the set of these collections by $\mathcal{C}$. $G$ has a subgraph on $k$ vertices with at least $z$ edges if and only if $\mathcal{C}$ contains a subset of $E(G)$. The lemma holds if this happens with probability smaller than

$\frac{1}{2}$. Since each member of $\mathcal{C}$ is a subset of $E(G)$ with probability $p^z$, the lemma follows from the inequality $|\mathcal{C}|p^z < \frac{1}{2}$,which can be shown from Lemma 5 as follows:

$$\begin{aligned}
|\mathcal{C}|p^z &\leq \binom{n}{k}\binom{\binom{k}{2}}{z}p^z \\
&< \left(\frac{en}{k}\right)^k \left(\frac{e\binom{k}{2}}{z}\right)^z p^z \\
&\leq \left(\frac{en}{k}\right)^k \left(\frac{e\binom{k}{2}}{\frac{8e^2\cdot\log(en/k)}{k\log(em/k)}\binom{k}{2}} \cdot \frac{4}{m-1}\log(en/m)\right)^z \\
&= \left(\frac{en}{k}\right)^k \left(\frac{1}{2e}\cdot\frac{k}{m-1}\cdot\frac{\log(en/m)}{\log(en/k)}\cdot\log(em/k)\right)^z \\
&< \left(\frac{en}{k}\right)^k \left(\frac{1}{2e}\cdot\frac{k}{m/2}\cdot 1\cdot\sqrt{em/k}\right)^z \\
&= \left(\frac{en}{k}\right)^k \left(\frac{k}{em}\right)^{\frac{z}{2}} \\
&= e^{k\log(en/k)-\frac{z}{2}\log(em/k)} \\
&< \frac{1}{2}.
\end{aligned}$$

□

## 2.2 The case $k \geq m$

It follows from the special case $m = k$ in Section 2.1 (and was also shown in [Al96]) that for any $s \geq m$ there is a graph $G(s,m)$ on $s$ vertices with $\alpha(G(s,m)) \leq m$ having no subgraph on $m$ vertices of density at least $8e^2\frac{\log(es/m)}{m}$. Let $n \geq k \geq m \geq 2$. Put $s = \frac{nm}{k}$ and $t = \frac{k}{m}$. For simplicity, assume that $s$ and $t$ are integers (using roundings one can get the proof in a similar way also in other cases). Set $G_0 = G(s,m)$. Replace each vertex in $G_0$ by $K_t$, and denote the obtained graph by $G$, i.e., $G = (V,E)$, where

$$\begin{aligned}
V &= V(G_0)\times\{1,2,\ldots,t\}, \\
E &= \left\{\{(v_1,i_1),(v_2,i_2)\} \in \binom{V}{2} : v_1 = v_2 \text{ or } (v_1,v_2)\in E(G_0)\right\}.
\end{aligned}$$

We now prove that $G$ gives the upper bound in Theorem 1 in case $k \geq m$. Certainly, $|V| = n$ and $\alpha(G) = \alpha(G_0) \leq m$. It remains to show that $G$ has no subgraph on $k$ vertices of density at least $c\cdot\frac{\log(en/k)}{m}$, which we now show with $c = 8e^2+1$.

We say that a subgraph $H \subseteq G$ *splits* a vertex $v \in V(G_0)$ if there are two indices $i,j \in \{1,2,\ldots,t\}$ with $(v,i) \in V(H), (v,j) \notin V(H)$. Let $H$ be a subgraph of $G$ on $k$

vertices with the maximum density (i.e., with the maximum number of edges). If $H$ splits two vertices $v_1, v_2 \in V(G_0)$, then we may find another (induced) subgraph $H' \subseteq G$ on $k$ vertices which splits fewer vertices of $G_0$ than $H$ and has density at least as big as $H$ ($H'$ can be obtained from $H$ by replacing either some vertices $(v_1, i)$ by some vertices $(v_2, j)$ or some vertices $(v_2, i)$ by some vertices $(v_1, j)$ in $H$). Repeating this procedure, we get a subgraph of $G$ on $k$ vertices which splits at most one vertex of $G_0$ and has density at least as big as $H$. Therefore, we may assume that $H$ itself splits at most one vertex of $G_0$. $H$ cannot split exactly one vertex of $G_0$, since the number of vertices in $H$ is divisible by $t$. Thus, we may assume that $H$ splits no vertex of $G_0$. By the construction of $G$, the density of $H$ is then smaller than

$$8e^2\frac{\log(es/m)}{m} + \frac{m\binom{t}{2}}{\binom{mt}{2}} < 8e^2\frac{\log(en/k)}{m} + \frac{1}{m} \leq (8e^2+1)\frac{\log(en/k)}{m}.$$

# 3 The lower bound

## 3.1 The case $m \geq k$

We now settle the most complicated case in the paper. The main idea (contained in Lemma 7(a) below) has some similarities with the ideas of Rödl [Rö86] and Thomason [Th87, Th88] leading to the currently asymptotically best upper bounds on the Ramsey off-diagonal numbers $r(k, m)$. These ideas are described in a simple way in [Ne95, pp. 1348-1349].

For $\sigma \in [0, 1]$, $m \geq 1$, $k \geq 1$, we define a Ramsey-type number $n_\sigma(m, k)$ as the minimum number $n$ such that any graph on $n$ vertices with independence number at most $m$ has a subgraph on $k$ vertices of density at least $\sigma$. (The density of a graph with one vertex is assumed to be 1.)

**Observation 6** *If $n_\sigma(m, k) \leq n$, then $\sigma(n, m, k) \geq \sigma$.*

Hence, an upper bound on the function $n_\sigma$ gives a lower bound on the function $\sigma$. Here is the key lemma:

**Lemma 7** *Let $\sigma \in [0, \frac{1}{2}]$ and let $\frac{1}{2\sigma}$ be an integer. Then*

*(a)* $n_\sigma(m, k) < n_\sigma(m - \frac{1}{2\sigma}, k) + n_\sigma(m, k - \frac{1}{2\sigma}) + \frac{1}{2\sigma}$, *for* $m \geq \frac{1}{\sigma}$ *and* $k \geq \frac{3}{4\sigma}$,

*(b)* $n_\sigma(\frac{1}{2\sigma}, k) = k$, *for* $k \geq \frac{1}{\sigma}$,

*(c)* $n_\sigma(m, \frac{1}{2\sigma}) \leq m + \frac{1}{8\sigma}$, *for* $m \geq \frac{1}{2\sigma}$.

*Proof.* **(a)** Suppose for a contrary that $G$ is a graph on $n_\sigma(m - \frac{1}{2\sigma}, k) + n_\sigma(m, k - \frac{1}{2\sigma}) + \frac{1}{2\sigma} - 1$ vertices with $\alpha(G) \leq m$ containing no subgraph on $k$ vertices of density at least $\sigma$. Since $G$ has more than $n_\sigma(m - \frac{1}{2\sigma}, k)$ vertices, it has $m - \frac{1}{2\sigma} \geq \frac{1}{2\sigma}$ independent vertices. Fix any set $I$ of $\frac{1}{2\sigma}$ independent vertices in $G$. Partition the set $V(G) \setminus I$ into two subsets $I_0$ and $I_1$ containing

the vertices of $V(G)\setminus I$ adjacent to no vertex in $I$ and to at least one vertex in $I$, respectively. By the pigeonhole principle, $|I_0| \geq n_\sigma(m-\frac{1}{2\sigma},k)$ or $|I_1| \geq n_\sigma(m,k-\frac{1}{2\sigma})$. In the first case $I_0$ contains $m-\frac{1}{2\sigma}$ independent vertices (otherwise $G$ would have a subgraph on $k$ vertices of density at least $\sigma$), which form, together with the vertices of $I$, $m$ independent vertices in $G$ – a contradiction. If $|I_1| \geq n_\sigma(m,k-\frac{1}{2\sigma})$, then there is a vertex set $I_2 \subseteq I_1, |I_2| = k-\frac{1}{2\sigma}$, such that the density of the subgraph of $G$ induced by $I_2$ is at least $\sigma$. Since

$$\left|\binom{I}{2}\right| = \frac{1}{2\sigma}(\frac{1}{4\sigma}-\frac{1}{2}) < \frac{1}{2\sigma}(k-\frac{1}{2\sigma}) = \frac{1}{2\sigma}|I_2|,$$

the density of the subgraph induced by the $k$ vertices of $I \cup I_2$ is then

$$\frac{\left|\binom{I_2}{2} \cap E(G)\right| + |(I\times I_2)\cap E(G)|}{\left|\binom{I}{2}\right| + \left|\binom{I_2}{2}\right| + |I\times I_2|} > \frac{\sigma\left|\binom{I_2}{2}\right| + |I_2|}{\frac{1}{2\sigma}|I_2| + \left|\binom{I_2}{2}\right| + \frac{1}{2\sigma}|I_2|} = \sigma,$$

which is again a contradiction. This completes the proof of (a).

**(b)** Let $G$ be a graph on $k \geq \frac{1}{\sigma}$ vertices with independence number at most $\frac{1}{2\sigma}$. By Turán's theorem [Tu54], the density of $G$ is at least

$$\frac{\frac{1}{2\sigma}\binom{2\sigma k}{2}}{\binom{k}{2}} = \frac{2\sigma k-1}{k-1} = \frac{\sigma k}{k-1} + \frac{\sigma k-1}{k-1} > \sigma + 0 = \sigma.$$

By Observation 3, $G$ contains a subgraph on $k$ vertices of density at least $\sigma$.

**(c)** Let $G$ be a graph on $m+\frac{1}{8\sigma}$ vertices with $\alpha(G) \leq m$. Certainly, $G$ contains at least $\frac{1}{8\sigma}$ edges. The density of any subgraph of $G$ on $\frac{1}{2\sigma}$ vertices with at least $\frac{1}{8\sigma}$ edges is at least

$$\frac{\frac{1}{8\sigma}}{\frac{\frac{1}{2\sigma}(\frac{1}{2\sigma}-1)}{2}} > \sigma.$$

□

**Consequence 8** *If $\frac{1}{2\sigma}$ is an integer and $m$, $k$ are two positive integers divisible by $\frac{1}{2\sigma}$, then*

$$n_\sigma(m,k) \;<\; \frac{1}{\sigma}\binom{2\sigma(m+k)}{2\sigma k} - \frac{1}{2\sigma}.$$

*Proof.* We fix $\sigma$ such that $\frac{1}{2\sigma}$ is an integer, and proceed by induction on $m+k$.

For $k = \frac{1}{2\sigma}$, Consequence 8 follows from Lemma 7(c):

$$n_\sigma(m,\frac{1}{2\sigma}) \leq m + \frac{1}{8\sigma} < \frac{1}{\sigma}(2\sigma m+1) - \frac{1}{2\sigma} = \frac{1}{\sigma}\binom{2\sigma(m+k)}{2\sigma k} - \frac{1}{2\sigma}.$$

For $m = \frac{1}{2\sigma}$ and $k > \frac{1}{2\sigma}$, Consequence 8 follows from Lemma 7(b):

$$n_\sigma(\frac{1}{2\sigma}, k) = k < \frac{1}{\sigma}(2\sigma k + 1) - \frac{1}{2\sigma} = \frac{1}{\sigma}\binom{2\sigma(m+k)}{2\sigma k} - \frac{1}{2\sigma}.$$

To complete the proof, it suffices to exhibit the second inductive step for $m > \frac{1}{2\sigma}, k > \frac{1}{2\sigma}$:

$$\begin{aligned} n_\sigma(m,k) &< n_\sigma(m - \frac{1}{2\sigma}, k) + n_\sigma(m, k - \frac{1}{2\sigma}) + \frac{1}{2\sigma} \\ &< \frac{1}{\sigma}\binom{2\sigma(m - \frac{1}{2\sigma} + k)}{2\sigma k} - \frac{1}{2\sigma} + \frac{1}{\sigma}\binom{2\sigma(m + k - \frac{1}{2\sigma})}{2\sigma(k - \frac{1}{2\sigma})} - \frac{1}{2\sigma} + \frac{1}{2\sigma} \\ &= \frac{1}{\sigma}\left(\binom{2\sigma(m+k)-1}{2\sigma k} + \binom{2\sigma(m+k)-1}{2\sigma k - 1}\right) - \frac{1}{2\sigma} \\ &= \frac{1}{\sigma}\binom{2\sigma(m+k)}{2\sigma k} - \frac{1}{2\sigma}. \end{aligned}$$

□

We can now prove the lower bound in Theorem 1 for $m \geq k$. We assume $m \geq k$ and set

$$\sigma_0 = \frac{\log(en/k)}{32k\log(em/k)}.$$

We distinguish three cases such that at least one of them must occur.
**Case 1:** $k \geq \frac{1}{2\sigma_0}$, $\sigma_0 \leq \frac{1}{2}$. This is the basic and most difficult case. Set

$$\sigma_1 = \frac{1}{2\left\lfloor \frac{1}{2\sigma_0} \right\rfloor}, \quad m_1 = \frac{\lceil 2\sigma_1 m \rceil}{2\sigma_1}, \quad k_1 = \frac{\lceil 2\sigma_1 k \rceil}{2\sigma_1}.$$

Obviously,

$$0 < \sigma_1 \leq \frac{1}{2}, \quad \sigma_0 \leq \sigma_1 < 2\sigma_0, \quad m \leq m_1 < 2m, \quad k \leq k_1 < 2k.$$

From Observation 3, Consequence 8, Lemma 5, and from $m \geq k \geq \frac{1}{2\sigma_0}$, we get

$$\begin{aligned} n_{\sigma_0}(m,k) &\leq n_{\sigma_1}(m_1, k_1) \\ &< \frac{1}{\sigma_1}\binom{2\sigma_1(m_1+k_1)}{2\sigma_1 k_1} \\ &< \frac{1}{\sigma_0}\binom{8\sigma_0(m+k)}{8\sigma_0 k} \\ &< \frac{1}{\sigma_0}\left(\frac{e(m+k)}{k}\right)^{8\sigma_0 k} \\ &\leq 2k\left(\frac{2em}{k}\right)^{\frac{\log(en/k)}{4\log(em/k)}} \end{aligned}$$

$$\begin{aligned} &= 2k\left(\frac{en}{k}\right)^{\frac{\log(2em/k)}{4\log(em/k)}} \\ &< 2k\left(\frac{en}{k}\right)^{\frac{1}{2}} \\ &= (4ekn)^{\frac{1}{2}} \\ &< n \end{aligned}$$

(the last inequality follows from $4ekn < \frac{kn}{e} \cdot e^{16} \leq \frac{kn}{e} \cdot e^{32k\sigma_0} = \frac{kn}{e} \cdot e^{\frac{\log(en/k)}{\log(em/k)}} \leq \frac{kn}{e} \cdot \frac{en}{k} = n^2$). From Observation 6 we now get

$$\boxed{\sigma(n,m,k) \geq \sigma_0 \quad \text{in Case 1.}}$$

**Case 2:** $\sigma_0 > \frac{1}{2}$**.** Consequence 8 and Lemma 5 give

$$\begin{aligned} n_{\frac{1}{2}}(m,k) &< 2\binom{m+k}{k} - 1 \\ &< 2\left(\frac{e(m+k)}{k}\right)^{k} - 1 \\ &< 2\left(\frac{2em}{k}\right)^{2\sigma_0 \cdot k} - 1 \\ &= 2\left(\frac{2em}{k}\right)^{\frac{\log(en/k)}{16\log(em/k)}} - 1 \\ &= 2\left(\frac{en}{k}\right)^{\frac{\log(2em/k)}{16\log(em/k)}} - 1 \\ &< 2\left(\frac{en}{k}\right)^{\frac{1}{8}} - 1 \\ &< n. \end{aligned}$$

Observation 6 now gives

$$\boxed{\sigma(n,m,k) \geq \tfrac{1}{2} \quad \text{in Case 2.}}$$

**Case 3:** $k < \frac{1}{2\sigma_0}$**.** Any graph on $n$ vertices with $\alpha(G) \leq m$ has at least $n - m \geq m \geq k$ edges (we have $m \leq \frac{n}{2}$), and thus any subgraph on $k$ vertices with $\lfloor k/2 \rfloor$ edges has density at least

$$\frac{\left\lfloor \frac{k}{2} \right\rfloor}{\frac{k(k-1)}{2}} > \frac{\frac{k}{4}}{\frac{k^2}{2}} = \frac{1}{2k} > \sigma_0.$$

Thus,

$$\boxed{\sigma(n,m,k) > \sigma_0 \quad \text{in Case 3.}}$$

We conclude that

$$\boxed{\sigma(n,m,k) \geq \min\left\{\sigma_0, \tfrac{1}{2}\right\} \quad \text{in each case.}}$$

Thus,

$$\sigma(n,m,k) \geq \min\left\{\sigma_0, \frac{1}{2}\right\} \geq \frac{1}{32}\min\left\{\frac{\log(en/k)}{k\log(em/k)}, 1\right\},$$

which is a desired lower bound in Theorem 1 for $m \geq k$.

## 3.2 The case $k \geq m$

Let $k \geq m$. We first suppose that $k \leq \frac{n}{e^3}$. We may assume that $k \geq 4$ (the case $k \leq 3$ is trivial). For a contrary, suppose that $G$ is a graph on $n$ vertices with $\alpha(G) \leq m$ such that the density of any subgraph on $k$ vertices is smaller than $\sigma = \frac{1}{4}\min\{\frac{\log(en/k)}{m}, 1\} = \min\{\frac{\log(en/k)}{4m}, \frac{1}{4}\}$. By Observation 3, every subgraph of $G$ on at least $k$ vertices has density smaller than $\sigma$.

We now find $m+1$ independent vertices $v_1, \ldots, v_{m+1}$, which gives a contradiction to $\alpha(G) \leq m$. We start with the graph $G$, and proceed inductively in $m+1$ steps $S_1, \ldots, S_{m+1}$ so that in step $S_i$ we choose a vertex $v_i$ of minimum degree in the current graph and remove it and all its neighbors from the graph. Let $V_i$ be the set of vertices which remain after step $S_i$. Thus, if $V_0$ denotes the vertex set of $G$, then $V_i = V_{i-1} \setminus (\{v_i\} \cup N_{i-1}(v_i))$, where $v_i$ is a vertex of minimum degree in the subgraph $G_{i-1}$ of $G$ induced by $V_{i-1}$ and $N_{i-1}(v_i)$ is the set of its neighbors in $G_{i-1}$. Certainly, the vertices $v_1, \ldots, v_{m+1}$ are independent. To prove the correctness of the argument, it suffices to show that $|V_i| > 0$ for $i = 1, \ldots, m$.

Let $i \in \{1, \ldots, m\}$. If $|V_{i-1}| \geq k$, then the density of $G_{i-1}$ is smaller than $\sigma$, and therefore the degree of $v_i$ in $G_{i-1}$ is smaller than $\sigma(|V_{i-1}| - 1)$. From this we get

$$|V_i| > |V_{i-1}| - 1 - \sigma(|V_{i-1}| - 1) > (1 - 2\sigma)|V_{i-1}|,$$

provided $|V_{i-1}| \geq k$ (since $\sigma|V_{i-1}| \geq 1$ follows from $|V_{i-1}| \geq k$, $\sigma = \min\{\frac{\log(en/k)}{4m}, \frac{1}{4}\}$, $k \geq 4$, and from our supposition $k \leq \frac{n}{e^3}$). Further, we get

$$|V_i| \;>\; (1 - 2\sigma)^i |V_0|,$$

provided $|V_{i-1}| \geq k$. Since the function $f(t) = 1 - 2t - e^{-3t}$ is concave and its values in $t = 0$ and $t = \frac{1}{4}$ are non-negative, we have $f(\sigma) \geq 0$, thus

$$1 - 2\sigma \geq e^{-3\sigma}.$$

Thus,

$$\begin{aligned} |V_i| &> (1 - 2\sigma)^i |V_0| \\ &\geq (1 - 2\sigma)^m |V_0| \end{aligned}$$

$$\begin{aligned} &\geq (e^{-3\sigma})^m n \\ &\geq e^{-\frac{3\log(en/k)}{4}} n^{\frac{3}{4}} n^{\frac{1}{4}} \\ &\geq \left(\frac{en}{k}\right)^{-\frac{3}{4}} n^{\frac{3}{4}} (e^3 k)^{\frac{1}{4}} \\ &= k, \end{aligned}$$

provided $|V_{i-1}| \geq k$. From this and from $|V_0| = n \geq k$, we get by induction on $i$ that

$$|V_1| \geq |V_2| \geq \cdots \geq |V_m| \geq k > 0,$$

which completes the argument for $k \leq \frac{n}{e^3}$.

If $k > \frac{n}{e^3}$ (and $m \leq \frac{n}{2}$) then, by Turán's theorem [Tu54], any graph $G$ on $n$ vertices with $\alpha(G) \leq m$ has density at least

$$\frac{\frac{n\left(\frac{n}{m}-1\right)}{2}}{\binom{n}{2}} > \frac{\frac{n\frac{n}{2m}}{2}}{\frac{n^2}{2}} = \frac{1}{2m} > \frac{\log(en/k)}{8m},$$

and the lower bound in Theorem 1 follows from Observation 3.

**Acknowledgments.** I would like to thank Noga Alon for interesting discussions and for helpful comments on the draft of this paper. I also thank the referee for helpful comments and suggestions.

# References

[Al96] N. Alon, Independence numbers of locally sparse graphs and a Ramsey type problem, Random Structures and Algorithms 9 (1996), 271–278.

[Bo85] B. Bollobás, Random graphs, Academic Press, London 1985.

[Er79] P. Erdős, Some old and new problems in various branches of Combinatorics, in: Proc. $10^{th}$ Southeastern Conf. on Combinatorics, Graph Theory and Computing, Boca Raton 1979 Vol I, Congressus Numerantium 23 (1979), 19-37.

[GS95] E. Győri and V.T. Sós (eds.), Problem collection of the DIMANET Mátraháza Workshop, Mátraháza, October 22–28, 1995.

[Ne95] J. Nešetřil, Ramsey theory, in: R.L. Graham, M. Grötschel, and L. Lovász (eds.), Handbook of combinatorics, Vol. II, Elsevier, New York 1995, pp. 1331–1403.

[Rö86] V. Rödl, Upper bounds on Ramsey numbers $R(k, l)$, unpublished manuscript (1986).

[Th87] A. Thomason, Random graphs, strongly regular graphs and pseudorandom graphs, in: C. Whitehead (ed.), Survey in combinatorics, Cambridge Univesity Press, Cambridge 1987, 173-196.

[Th88] A. Thomason, An upper bound for some Ramsey numbers, J. Graph Theory 12 (1988), 509–517.

[Tu54] P. Turán, On the theory of graphs, Colloquium Math. 3 (1954), 19–30.