DIMACS Technical Report 99-37
June 1999

New bounds on crossing numbers

by

János Pach[1]
Courant Institute, NYU and Hungarian Academy of Sciences

Joel Spencer
Courant Institute, NYU

Géza Tóth[2]
Massachusetts Institute of Technology
and Hungarian Academy of Sciences

[1]Supported by NSF grant CCR-97-32101 and PSC-CUNY Research Award 667339.
[2]Supported by DIMACS Center, OTKA-T-020914, and OTKA-F-22234.

DIMACS is a partnership of Rutgers University, Princeton University, AT&T Labs-Research, Bell Labs, Telcordia Technologies (formerly Bellcore) and NEC Research Institute.

DIMACS is an NSF Science and Technology Center, funded under contract STC–91–19999; and also receives support from the New Jersey Commission on Science and Technology.

ABSTRACT

The *crossing number*, $\mathrm{cr}(G)$, of a graph G is the least number of crossing points in any drawing of G in the plane. Denote by $\kappa(n, e)$ the minimum of $\mathrm{cr}(G)$ taken over all graphs with n vertices and at least e edges. We prove a conjecture of P. Erdős and R. Guy by showing that $\kappa(n, e)n^2/e^3$ tends to a positive constant as $n \to \infty$ and $n \ll e \ll n^2$. Similar results hold for graph drawings on any other surface of fixed genus.

We prove better bounds for graphs satisfying some monotone properties. In particular, we show that if G is a graph with n vertices and $e \geq 4n$ edges, which does not contain a cycle of length *four* (resp. *six*), then its crossing number is at least ce^4/n^3 (resp. ce^5/n^4), where $c > 0$ is a suitable constant. These results cannot be improved, apart from the value of the constant. This settles a question of M. Simonovits.

1 Introduction

Let G be a simple undirected graph with $n(G)$ nodes (vertices) and $e(G)$ edges. A *drawing* of G in the *plane* is a mapping f that assigns to each vertex of G a distinct point in the plane and to each edge uv a continuous arc connecting $f(u)$ and $f(v)$, not passing through the image of any other vertex. For simplicity, the arc assigned to uv is also called an *edge*, and if this leads to no confusion, it is also denoted by uv. We assume that no three edges have an interior point in common. The *crossing number*, $\mathrm{cr}(G)$, of G is the minimum number of crossing points in any drawing of G.

The determination of $\mathrm{cr}(G)$ is an NP-complete problem [GJ83]. It was discovered by Leighton [L84] that the crossing number can be used to estimate the chip area required for the VLSI circuit layout of a graph. He proved the following general lower bound for $\mathrm{cr}(G)$, which was discovered independently by Ajtai, Chvátal, Newborn, and Szemerédi. The best known constant, 1/33.75, in the theorem is due to Pach and Tóth.

Theorem A. [ACNS82], [L84], [PT97] *Let G be a graph with $n(G) = n$ nodes and $e(G) = e$ edges, $e \geq 7.5n$. Then we have*

$$\mathrm{cr}(G) \geq \frac{1}{33.75}\frac{e^3}{n^2}.$$

Theorem A can be used to deduce the best known upper bounds for the number of unit distances determined by n points in the plane [S98], for the number of different ways how a line can split a set of n points into two equal parts [D98], and it has some other interesting corollaries [PS98].

It is easy to see that the bound in Theorem A is tight, apart from the value of the constant. However, as it was suggested by Miklós Simonovits [S97], it may be possible to strengthen the theorem for some special classes of graphs, e.g., for graphs not containing some fixed, so-called *forbidden* subgraph. In Sections 2 and 3 of the present paper we verify this conjecture.

A graph property $\mathcal{P}$ is said to be *monotone* if

- if a graph G satisfies $\mathcal{P}$, then every subgraph of G also satisfies $\mathcal{P}$;
- if G_1 and G_2 satisfy $\mathcal{P}$, then their disjoint union also satisfies $\mathcal{P}$.

For any monotone property $\mathcal{P}$, let $\mathrm{ex}(n, \mathcal{P})$ denote the maximum number of edges that a graph of n vertices can have if it satisfies $\mathcal{P}$. In the special case when $\mathcal{P}$ is the property that the graph does not contain a subgraph isomorphic to a fixed forbidden subgraph H, we write $\mathrm{ex}(n, H)$ for $\mathrm{ex}(n, \mathcal{P})$.

Theorem 1. *Let $\mathcal{P}$ be a monotone graph property with $\mathrm{ex}(n, \mathcal{P}) = O(n^{1+\alpha})$ for some $\alpha > 0$.*

Then there exist two constants $c, c' > 0$ such that the crossing number of any graph G with property $\mathcal{P}$, which has n vertices and $e \geq cn\log^2 n$ edges, satisfies

$$\mathrm{cr}(G) \geq c'\frac{e^{2+1/\alpha}}{n^{1+1/\alpha}}.$$

If $\mathrm{ex}(n, \mathcal{P}) = \Theta(n^{1+\alpha})$*, then this bound is asymptotically tight, up to a constant factor.*

In some interesting special cases when we know the precise order of magnitude of the function $\mathrm{ex}(n, \mathcal{P})$, we obtain some slightly stronger results. The *girth* of a graph is the length of its shortest cycle.

Theorem 2. *Let G be a graph with n vertices and $e \geq 4n$ edges, whose girth is larger than $2r$, for some $r > 0$ integer. Then the crossing number of G satisfies*

$$\mathrm{cr}(G) \geq c_r \frac{e^{r+2}}{n^{r+1}},$$

where $c_r > 0$ is a suitable constant. For $r = 2, 3,$ and 5, these bounds are asymptotically tight, up to a constant factor.

What happens if the girth of G is larger than $2r + 1$? Since one can destroy every odd cycle of a graph by deleting at most half of its edges, even in this case we cannot expect an asymptotically better lower bound for the crossing number of G than the bound given in Theorem 2.

Theorem 3. *Let G be a graph with n vertices and $e \geq 4n$ edges, which does not contain a complete bipartite subgraph $K_{r,s}$ with r and s vertices in its classes, $s \geq r$.*

Then the crossing number of G satisfies

$$\mathrm{cr}(G) \geq c_{r,s} \frac{e^{3+1/(r-1)}}{n^{2+1/(r-1)}},$$

where $c_{r,s} > 0$ is a suitable constant. These bounds are tight up to a constant factor if $r = 2, 3,$ or if r is arbitrary and $s > (r-1)!$.

The *bisection width*, $b(G)$, of a graph G is defined as the minimum number of edges whose removal splits the graph into two roughly equal subgraphs. More precisely, $b(G)$ is the minimum number of edges running between V_1 and V_2, over all partitions of the vertex set of G into two parts $V_1 \cup V_2$ such that $|V_1|, |V_2| \geq n(G)/3$.

Leighton [L83] observed that there is an intimate relationship between the bisection width and the crossing number of a graph, which is based on the Lipton–Tarjan separator theorem for planar graphs [LT79]. The proofs of Theorems 1-3 are based on repeated application of the following version of this relationship.

Theorem B. [PSS96] *Let G be a graph of n vertices, whose degrees are $d_1, d_2, \ldots, d_n$. Then*

$$b(G) \leq 10\sqrt{\mathrm{cr}(G)} + 2\sqrt{\sum_{i=1}^{n} d_i^2}.$$

Let $\kappa(n, e)$ denote the minimum crossing number of a graph G with n vertices and at least e edges. That is,

$$\kappa(n, e) = \min_{\substack{n(G) = n \\ e(G) \geq e}} \mathrm{cr}(G).$$

It follows from Theorem A that, for $e \geq 4n$, $\kappa(n,e)n^2/e^3$ is bounded from below and from above by two positive constants. Paul Erdős and Richard K. Guy [EG73] conjectured that if $e \gg n$ then $\lim \kappa(n,e)n^2/e^3$ exists. (We use the notation $f(n) \gg g(n)$ to express that $\lim_{n\to\infty} f(n)/g(n) = \infty$.) In Section 4, we settle this problem.

Theorem 4. *If $n \ll e \ll n^2$, then*

$$\lim_{n\to\infty} \kappa(n,e)\frac{n^2}{e^3} = C > 0$$

exists.

We call the constant $C > 0$ in Theorem 4 the *midrange crossing constant*. It is necessary to limit the range of e from below and from above. (See the Remark at the end of Section 4.)

All of the above problems can be reformulated for graph drawings on other surfaces. Let S_g denote a torus with g holes, i.e., a compact oriented surface of *genus* g with no boundary. Define $\mathrm{cr}_g(G)$, the crossing number of G on S_g, as the minimum number of crossing points in any drawing of G on S_g. Let

$$\kappa_g(n,e) = \min_{\substack{n(G)=n \\ e(G)\geq e}} \mathrm{cr}_g(G).$$

With this notation, $\mathrm{cr}_0(G)$ is the planar crossing number and $\kappa_0(n,e) = \kappa(n,e)$.

In Section 5, we prove that there is a midrange crossing constant for graph drawings on any surface S_g of fixed genus $g \geq 0$.

Theorem 5. *For every $g \geq 0$, if $n \ll e \ll n^2$ then the limit*

$$\lim_{n\to\infty} \kappa_g(n,e)\frac{n^2}{e^3}$$

exists and is equal to the constant $C > 0$ in Theorem 4.

To prove this result, we have to generalize Theorem B.

Theorem 6. *Let G be a graph with n vertices, whose degrees are $d_1, d_2, \ldots, d_n$. Then*

$$b(G) \leq 300(1+g^{3/4})\sqrt{\mathrm{cr}_g(G) + \sum_{i=1}^{n} d_i^2}.$$

2 Crossing numbers and monotone properties – Proof of Theorem 1

Let $\mathcal{P}$ be a monotone graph property with $\mathrm{ex}(n,\mathcal{P}) \leq An^{1+\alpha}$, for some $A, \alpha > 0$. Let G be a graph with vertex set $V(G)$ and edge set $E(G)$, where $|V(G)| = n(G) = n$ and

$|E(G)| = e(G) = e$. Suppose that G satisfies property $\mathcal{P}$ and $e \geq cn\log^2 n$. To prove Theorem 1, we assume that

$$\mathrm{cr}(G) < c'\frac{e^{2+1/\alpha}}{n^{1+1/\alpha}},$$

and, if c and c' are suitable constant, we will obtain a contradiction.

We break G into smaller components, according to the following procedure.

Decomposition Algorithm

Step 0. **Let** $G^0 = G, G_1^0 = G, M_0 = 1, m_0 = 1$.

Suppose that we have already executed Step i, and that the resulting graph, G^i, consists of M_i components, $G_1^i, G_2^i, \ldots, G_{M_i}^i$, each of at most $(2/3)^i n$ vertices. Assume, without loss of generality, that the first m_i components of G^i have at least $(2/3)^{i+1}n$ vertices and the remaining $M_i - m_i$ have fewer. Then

$$(2/3)^{i+1}n(G) \leq n(G_j^i) \leq (2/3)^i n(G) \quad (j = 1, 2, \ldots, m_i).$$

Thus, we have that $m_i \leq (3/2)^{i+1}$.

Step $i+1$. **If**

$$(2/3)^i < \frac{1}{(2A)^{1/\alpha}} \cdot \frac{e^{1/\alpha}}{n^{1+1/\alpha}}, \tag{1}$$

then stop. (1) is called the *stopping rule*.

Else, for $j = 1, 2, \ldots, m_i$, delete $b(G_j^i)$ edges from G_j^i such that G_j^i falls into two components, each of at most $(2/3)n(G_j^i)$ vertices. Let G^{i+1} denote the resulting graph on the original set of n vertices. Clearly, each component of G^{i+1} has at most $(2/3)^{i+1}n$ vertices.

Suppose that the Decomposition Algorithm terminates in Step $k+1$. If $k > 0$, then

$$(2/3)^k < \frac{1}{(2A)^{1/\alpha}} \cdot \frac{e^{1/\alpha}}{n^{1+1/\alpha}} \leq (2/3)^{k-1}.$$

First, we give an upper bound on the total number of edges deleted from G.

Using that, for any non-negative reals $a_1, a_2, \ldots, a_m$,

$$\sum_{j=1}^{m} \sqrt{a_j} \leq \sqrt{m\sum_{j=1}^{m} a_j}, \tag{2}$$

we obtain that, for any $0 \leq i < k$,

$$\sum_{j=1}^{m_i} \sqrt{\mathrm{cr}(G_j^i)} \leq \sqrt{m_i \sum_{j=1}^{m_i} \mathrm{cr}(G_j^i)} \leq \sqrt{(3/2)^{i+1}}\sqrt{\mathrm{cr}(G)} < \sqrt{(3/2)^{i+1}}\sqrt{\frac{c'e^{2+1/\alpha}}{n^{1+1/\alpha}}}.$$

Denoting by $d(v, G_j^i)$ the degree of vertex v in G_j^i, we have

$$\sum_{j=1}^{m_i} \sqrt{\sum_{v\in V(G_j^i)} d^2(v, G_j^i)} \leq \sqrt{(3/2)^{i+1}}\sqrt{\sum_{v\in V(G^i)} d^2(v, G^i)}$$

$$\leq \sqrt{(3/2)^{i+1}} \sqrt{\max_{v \in V(G^i)} d(v, G^i) \sum_{v \in V(G^i)} d(v, G^i)} \leq \sqrt{(3/2)^{i+1}} \sqrt{(2/3)^i n(2e)} = \sqrt{3en}.$$

In view of Theorem B in the Introduction, the total number of edges deleted during the procedure is

$$\sum_{i=0}^{k-1} \sum_{j=1}^{m_i} b(G_j^i) \leq 10 \sum_{i=0}^{k-1} \sum_{j=1}^{m_i} \sqrt{\mathrm{cr}(G_j^i)} + 2 \sum_{i=0}^{k-1} \sum_{j=1}^{m_i} \sqrt{\sum_{v \in V(G_j^i)} d^2(v, G_j^i)}$$

$$< 10\sqrt{c'} \sqrt{\frac{e^{2+1/\alpha}}{n^{1+1/\alpha}}} \sum_{i=0}^{k-1} \sqrt{(3/2)^i} + 2k\sqrt{3en} \leq 250\sqrt{c'} \sqrt{\frac{e^{2+1/\alpha}}{n^{1+1/\alpha}}} \sqrt{(2A)^{1/\alpha} \frac{n^{1+1/\alpha}}{e^{1/\alpha}}} + 2k\sqrt{3en} \leq \frac{e}{2},$$

provided that c' is sufficiently small and c is sufficiently large.

Therefore, the number of edges of the graph G^k obtained in the final STEP of the algorithm satisfies

$$e(G^k) \geq \frac{e}{2}.$$

(Note that this inequality trivially holds if the algorithm terminates in the very first STEP, i.e., when $k = 0$.)

Next we give a lower bound on $e(G^k)$. The number of vertices of each connected component of G^k satisfies

$$n(G_j^k) \leq (2/3)^k n < \frac{1}{(2A)^{1/\alpha}} \cdot \frac{e^{1/\alpha}}{n^{1+1/\alpha}} n = \left(\frac{e}{2An}\right)^{1/\alpha} \quad (j = 1, 2, \ldots, M_k).$$

Since each G_j^k has property $\mathcal{P}$, it follows that

$$e(G_j^k) \leq An^{1+\alpha}(G_j^k) < An(G_j^k) \cdot \frac{e}{2An}.$$

Therefore, for the total number of edges of G_k, we have

$$e(G^k) = \sum_{j=1}^{M_k} e(G_j^k) < A\frac{e}{2An} \sum_{j=1}^{M_k} n(G_j^k) = \frac{e}{2},$$

the desired contradiction. This proves the bound of Theorem 1.

It remains to show that the bound is tight up to a constant factor. Suppose that $\mathrm{ex}(n, \mathcal{P}) \geq A' n^{1+\alpha}$ For every e ($cn < e \leq An^{1+\alpha}$), we construct a graph G of at most n vertices and at least e edges, which has property $\mathcal{P}$ and crossing number

$$\mathrm{cr}(G) \leq c'' \frac{e^{2+1/\alpha}}{n^{1+1/\alpha}},$$

for a suitable constant $c'' = c''(A', \alpha)$.

Let

$$k = \left\lceil \frac{2e}{A'n} \right\rceil^{\frac{1}{\alpha}},$$

and let G_k denote a graph of k vertices and at least $A'k^{1+\alpha}$ edges, which has property $\mathcal{P}$. Clearly,

$$\mathrm{cr}(G_k) \le e^2(G_k) \le (Ak^{1+\alpha})^2 = A^2k^{2+2\alpha}.$$

Let G be the union of $\lfloor n/k \rfloor$ disjoint copies of G_k. Then $n(G) = \lfloor n/k \rfloor k \le n$,

$$e(G) = \left\lfloor \frac{n}{k} \right\rfloor e(G_k) \ge \frac{n}{2k} A'kk^{\alpha} \ge e,$$

$$\mathrm{cr}(G) = \left\lfloor \frac{n}{k} \right\rfloor \mathrm{cr}(G_k) \le \frac{n}{k} A^2k^{2+2\alpha} \le A^2n \left(2 \left(\frac{2e}{A'n} \right)^{\frac{1}{\alpha}} \right)^{1+2\alpha} = \frac{2^{3+2\alpha+1/\alpha}A^2}{(A')^{2+1/\alpha}} \cdot \frac{e^{2+1/\alpha}}{n^{1+1/\alpha}},$$

as required. □

3 Forbidden subgraphs – Proofs of Theorems 2 and 3

In Section 1, we established Theorem 1 under the assumption $e \ge cn\log^2 n$, where c is a suitable constant depending on property $\mathcal{P}$. It seems very likely that the same result is true for every $e \ge cn$. The appearance of the $\log^2 n$ factor was due to the fact that to estimate the total number of edges deleted during the Decomposition Algorithm, we applied Theorem B. We used a poor upper bound on the term $\sum d_i^2$, because some of the degrees d_i may be very large. However, in some interesting special cases, this difficulty can be avoided by a simple trick. We can split each vertex of high degree into vertices of 'average degree,' unless the new graph ceases to have property $\mathcal{P}$.

We illustrate this technique by proving the following result, which is the $r = s = 2$ special case of Theorem 3 and a slight modification of Theorem 2 for $r = 2$.

Theorem 3.1. *Let G be a $K_{2,2}$-free (C_4-free) graph with $n(G) = n$ vertices and $e(G) = e$ edges, $e \ge 1000n$. Then*

$$\mathrm{cr}(G) \ge \frac{1}{10^8} \frac{e^4}{n^3}.$$

This bound is tight up to a constant factor.

Proof. Let G be a graph with n vertices and $e \ge 1000n$ edges, which does not contain $K_{2,2}$ as a subgraph. Suppose, in order to obtain a contradiction, that

$$\mathrm{cr}(G) < \frac{1}{10^8} \frac{e^4}{n^3},$$

and G is drawn in the plane with $\mathrm{cr}(G)$ crossings.

First, we split every vertex of G whose degree exceeds $\overline{d} := 2e/n$ into vertices of degree at most $\overline{d}$, as follows. Let v be a vertex of G with degree $d(v, G) = d(v) = d > \overline{d}$, and let $vw_1, vw_2, \ldots, vw_d$ be the edges incident to v, listed in clockwise order. Replace v by $\lceil d/\overline{d} \rceil$ new vertices, $v_1, v_2, \ldots, v_{\lceil d/\overline{d} \rceil}$, placed in clockwise order on a very small circle around v. Without introducing any new crossings, connect w_j to v_i if and only if $\overline{d}(i-1) < j \leq \overline{d}i$ $(1 \leq j \leq d, 1 \leq i \leq \lceil d/\overline{d} \rceil)$. Repeat this procedure for every vertex whose degree exceeds $\overline{d}$, and denote the resulting graph by G'.

Obviously, G' is also $K_{2,2}$-free, $e(G') = e(G) = e$, and

$$\mathrm{cr}(G') \leq \mathrm{cr}(G) < \frac{1}{10^8} \frac{e^4(G)}{n^3(G)}.$$

Since all but at most n vertices of G' have degree $\overline{d}$, we have $n(G') < 2n(G) = 2n$.

Apply the Decomposition Algorithm described in the previous section to the graph G' with the difference that, instead of (1), use the following stopping rule: stop in Step $i+1$ if

$$\left(\frac{2}{3}\right)^i < \frac{e^2(G')}{16n^3(G')}.$$

Suppose that the algorithm terminates in Step $k+1$. If $k > 0$, then

$$(2/3)^k < \frac{e^2(G')}{16n^3(G')} \leq (2/3)^{k-1}.$$

Just like in the proof of Theorem 1, for every $i < k$, we have that

$$\sum_{j=1}^{m_i} \sqrt{\mathrm{cr}(G_j^i)} \leq \sqrt{(3/2)^{i+1}} \sqrt{\mathrm{cr}(G)} < \frac{1}{10^4} \sqrt{(3/2)^{i+1}} \frac{e^2}{n^{3/2}}$$

and, using the fact that the maximum degree in G' is at most $\overline{d}$,

$$\sum_{j=1}^{m_i} \sqrt{\sum_{v \in V(G_j^i)} d^2(v, G_j^i)} \leq \sqrt{(3/2)^{i+1}} \sqrt{\sum_{v \in V(G')} d^2(v, G')}$$

$$\leq \sqrt{(3/2)^{i+1}} \sqrt{\overline{d} 2e(G')} \leq 2\sqrt{(3/2)^{i+1}} \frac{e}{\sqrt{n}}.$$

Hence, by Theorem B, the total number of edges deleted during the algorithm is

$$\sum_{i=0}^{k-1} \sum_{j=1}^{m_i} b(G_j^i) \leq 10 \sum_{i=0}^{k-1} \sum_{j=1}^{m_i} \sqrt{\mathrm{cr}(G_j^i)} + 2 \sum_{i=0}^{k-1} \sum_{j=1}^{m_i} \sqrt{\sum_{v \in V(G_j^i)} d^2(v, G_j^i)}$$

$$< \frac{1}{1000} \frac{e^2}{n^{3/2}} \sum_{i=0}^{k-1} \sqrt{(3/2)^{i+1}} + 4 \frac{e}{\sqrt{n}} \sum_{i=0}^{k-1} \sqrt{(3/2)^{i+1}} = \sqrt{3/2} \frac{\sqrt{(3/2)^k} - 1}{\sqrt{3/2} - 1} \left(\frac{e^2}{1000 n^{3/2}} + \frac{4e}{\sqrt{n}} \right)$$

$$< 100\frac{n^{3/2}}{e}\left(\frac{e^2}{1000n^{3/2}} + \frac{4e}{\sqrt{n}}\right) < \frac{e}{10} + 400n < \frac{e}{2}.$$

Therefore, for the resulting graph,

$$e(G^k) \geq \frac{e}{2}.$$

On the other hand, each component of G^k has relatively few vertices:

$$n(G_j^k) < (2/3)^k n(G') < \frac{e^2}{16n^2(G')} = \frac{e^2}{16n^2(G^k)} \quad (j = 1, 2, \ldots, M_k).$$

Claim C. [R58] *Let* $\mathrm{ex}(n, K_{2,2})$ *denote the maximum number of edges that a* $K_{2,2}$*-free graph with* n *vertices can have. Then*

$$\mathrm{ex}(n, K_{2,2}) \leq \frac{n\left(1 + \sqrt{4n-3}\right)}{4} \leq n^{3/2}.$$

Applying the Claim to each G_k^j, we obtain

$$e(G_j^k) \leq n^{3/2}(G_j^k) < n(G_j^k) \cdot \sqrt{\frac{e^2}{16n^2(G^k)}},$$

therefore,

$$e(G^k) = \sum_{j=1}^{M_k} e(G_j^k) < \frac{e}{4n(G^k)} \sum_{j=1}^{M_k} n(G_j^k) = \frac{e}{4},$$

the desired contradiction. The tightness of Theorem 3.1 immediately follows from the fact that Theorem 1 was tight. □

Theorems 2 and 3 can be proved similarly. It is enough to notice that splitting a vertex of high degree does not decrease the girth of a graph G and does not create a subgraph isomorphic to $K_{r,s}$. Instead of Claim C, now we need

Claim C′. [BS74], [B66], [Be66], [S66], [W91] *For a fixed positive integer* r*, let* $\mathcal{G}_{2r}$ *denote the property that the girth of a graph is larger than* $2r$*.*

Then the maximum number of edges of a graph with n *vertices, which has property* $\mathcal{G}_{2r}$*, satisfies*

$$\mathrm{ex}(n, \mathcal{G}_{2r}) = O(n^{1+1/r}).$$

For $r = 2, 3$ *and* 5*, this bound is tight.*

Claim C″. [KST54], [F96], [ER62], [B66], [ARS98] *For any integers* $s \geq r \geq 2$*, the maximum number of edges of a* $K_{r,s}$*-free graph of* n *vertices, satisfies*

$$\mathrm{ex}(n, K_{r,s}) = O(n^{2-1/r}).$$

This bound is tight for $s > (r-1)!$*.*

In case $r = 3$, we obtain the following slight generalization of Theorem 2.

Theorem 3.2. *Let G be a graph of n vertices and $e \geq 4n$ edges, which contains no cycle C_6 of length 6.*

Then, for a suitable constant $c_6' > 0$, we have

$$\mathrm{cr}(G) \geq c_6' \frac{e^5}{n^4}.$$

To establish Theorem 3.2, it is enough to modify the proof of Theorem 2 at one point. Before splitting the high-degree vertices of G and running the Decomposition Algorithm, we have to turn G into a bipartite graph, by deleting at most half of its edges. After that, splitting a vertex cannot create a C_6, and the rest of the above argument shows that the crossing number of the remaining graph still exceeds $c_6' \frac{e^5}{n^4}$.

We do not see, however, how to obtain the analogous generalization of Theorem 2 for $r > 3$.

4 Midrange crossing constant in the plane – Proof of Theorem 4

Lemma 4.1. (i) *For any $a > 0$, the limit*

$$\gamma[a] = \lim_{n\to\infty} \frac{\kappa(n, na)}{n}$$

exists and is finite.
(ii) *$\gamma[a]$ is a convex continuous function.*
(iii) *For any $a \geq 4$, $1 > \delta > 0$,*

$$\gamma[a] - \gamma[a(1-\delta)] \leq \gamma[a(1+\delta)] - \gamma[a] \leq 10^3 \delta \gamma[a].$$

Proof. Clearly, any two graphs, G_1 and G_2, can be drawn in the plane so that the edges of G_1 do not intersect the edges of G_2. Therefore,

$$\kappa(n_1 + n_2, e_1 + e_2) \leq \kappa(n_1, e_1) + \kappa(n_2, e_2). \tag{3}$$

In particular, the function $f_a(n) = \kappa(n, na)$ is subadditive and hence the limit

$$\gamma[a] = \lim_{n\to\infty} \frac{\kappa(n, na)}{n}$$

exists and is finite for every fixed $a > 0$. It also follows from (3) that for any $a, b > 0$ and $1 > \alpha > 0$, if n and αn are both integers,

$$\kappa(n, (\alpha a + (1-\alpha)b)n) \leq \kappa(\alpha n, \alpha a n) + \kappa((1-\alpha)n, (1-\alpha)bn),$$

so for any $1 > \alpha > 0$ rational,

$$\gamma[\alpha a + (1-\alpha)b] \le \alpha\gamma[a] + (1-\alpha)\gamma[b].$$

But since the function $\gamma[a]$ is monotone increasing, it follows that for *any* $1 > \alpha > 0$,

$$\gamma[\alpha a + (1-\alpha)b] \le \alpha\gamma[a] + (1-\alpha)\gamma[b]. \tag{4}$$

That is, the function $\gamma[a]$ is *convex*. In particular, for every $1 > \delta > 0$, we have

$$\gamma[a] - \gamma[a(1-\delta)] \le \gamma[a(1+\delta)] - \gamma[a].$$

It is known that for any $a \ge 4$,

$$\frac{a^3 n}{100} \le \kappa(n, an) \le a^3 n \quad \Rightarrow \quad \frac{a^3}{100} \le \gamma[a] \le a^3 \tag{5}$$

(see e.g. [PT97]). Let $a \ge 4$, $1 > \delta > 0$. By (4),

$$\gamma[a(1+\delta)] \le (1-\delta)\gamma[a] + \delta\gamma[2a].$$

Therefore, using (5),

$$\gamma[a(1+\delta)] - \gamma[a] \le \delta\gamma[2a] \le \delta 8a^3 < 10^3\delta\gamma[a]. \ \square$$

Set

$$C := \limsup_{a\to\infty} \frac{\gamma[a]}{a^3}.$$

By (5), we have that $C < 1$.

Lemma 4.2. *For any $0 < \varepsilon < 1$, there exists $N = N(\varepsilon)$ such that $\kappa(n,e) > C\frac{\epsilon^3}{n^2}(1-\varepsilon)$, whenever $\min\{n, e/n, n^2/e\} > N$.*

Proof. Let $A > \frac{10^9}{\varepsilon^3}$ be a rational number satisfying

$$\frac{\gamma[A]}{A^3} > C(1 - \frac{\varepsilon}{10}). \tag{6}$$

Let $N = N(\varepsilon) \ge A$ such that, if $n > N$, $e = nA'$, and $|A - A'| \le A\varepsilon$, then

$$\kappa(n,e) > \gamma[A'](1 - \frac{\varepsilon}{10})n. \tag{7}$$

Let n and e be fixed, $\min\{n, e/n, n^2/e\} > N$ and let $G = (V, E)$ be a graph with $|V| = n$ vertices and $|E| = e$ edges, drawn in the plane with $\kappa(n,e)$ crossings. Set $p = An/e$. Let U be a randomly chosen subset of V with $\Pr[v \in U] = p$, independently for all $v \in V$. Let $\nu = |U|$, and let η (resp. ξ) be the number of edges (resp. crossings) in the (drawing of the) subgraph of G induced by the elements of U.

ν has mean pn and variance $p(1-p)n \le pn$, so, by the Chebyshev Inequality,

$$\Pr\left[|\nu - pn| > \frac{\varepsilon}{10^4}pn\right] < \frac{\varepsilon}{10}.$$

Write $\eta = \sum I_{uv}$, where the sum is taken over all edges $uv = vu \in E$, and I_{uv} denotes the indicator for the event $u, v \in U$. Obviously, $E[\eta] = \sum_{uv\in E} E[I_{uv}] = ep^2$. We decompose

$$\mathrm{Var}\left[\eta\right] = \sum_{uv\in E} \mathrm{Var}\left[I_{uv}\right] + \sum_{uv,uw\in E} Cov\left[I_{uv}, I_{uw}\right],$$

as $\mathrm{Cov}[I_{uv}, I_{wz}] = 0$ when all four indices are distinct. As always with indicators, we have

$$\sum_{uv\in E} \mathrm{Var}\left[I_{uv}\right] \le \sum_{uv\in E} E\left[I_{uv}\right] = E\left[\eta\right] = ep^2.$$

Using the bound $\mathrm{Cov}[I_{uv}, I_{uw}] \le E[I_{uv}I_{uw}] = p^3$, we obtain

$$\mathrm{Var}\left[\eta\right] \le p^2 e + p^3 \sum_{v\in V} \binom{d(v)}{2},$$

where $d(v)$ is the degree of vertex v in G. But $\sum_{v\in V} d(v) = 2e$ and all $d(v) < n$, so

$$\sum_{v\in V} \binom{d(v)}{2} \le \frac{1}{2}\sum_{v\in V} d^2(v) \le en.$$

Thus, we have

$$\mathrm{Var}\left[\eta\right] \le p^2 e + p^3 en \le 2p^3 en,$$

as $pn = An^2/e \ge 1$. Again, by the Chebyshev Inequality,

$$\Pr\left[|\eta - p^2 e| > \frac{\varepsilon}{10^4}p^2 e\right] < \frac{\varepsilon}{10}.$$

With probability at least $1 - \frac{\varepsilon}{5}$,

$$pn(1 - \frac{\varepsilon}{10^4}) < \nu < pn(1 + \frac{\varepsilon}{10^4}) \quad \text{and} \quad p^2 e(1 - \frac{\varepsilon}{10^4}) < \eta < p^2 e(1 + \frac{\varepsilon}{10^4}),$$

so with probability at least $1 - \frac{\varepsilon}{5}$,

$$A(1 - \frac{3\varepsilon}{10^4}) < \frac{\eta}{\nu} = A' < A(1 + \frac{3\varepsilon}{10^4}).$$

Therefore, in view of (7), with probability at least $1 - \frac{\varepsilon}{5}$, the subgraph of G induced by U has at least $pn(1 - \frac{\varepsilon}{10})\gamma[A'](1 - \frac{\varepsilon}{10})$ crossings. But then, we have

$$E\left[\xi\right] \ge (1 - \frac{\varepsilon}{5})pn(1 - \frac{\varepsilon}{10})\gamma[A'](1 - \frac{\varepsilon}{10}) \ge (1 - \frac{\varepsilon}{5})pn(1 - \frac{\varepsilon}{10})\gamma[A](1 - \frac{3\varepsilon}{10})(1 - \frac{\varepsilon}{10})$$

$$\geq (1-\frac{\varepsilon}{5})pn(1-\frac{\varepsilon}{10})CA^3(1-\frac{\varepsilon}{10})(1-\frac{3\varepsilon}{10})(1-\frac{\varepsilon}{10}) \geq (1-\varepsilon)CA^3pn,$$

where the second and third inequalities follow from Lemma 4.1 (iii) and from the choice of A, respectively.

On the other hand,

$$E[\xi] = p^4\kappa(n,e),$$

as every crossing lies in U with probability p^4. Thus

$$\kappa(n,e) \geq (1-\varepsilon)\frac{pnCA^3}{p^4} = C\frac{e^3}{n^2}(1-\varepsilon)$$

as desired. □

To complete the proof of Theorem 4, we have to establish the "counterpart" of Lemma 4.2.

Lemma 4.3. *For any $1 > \varepsilon > 0$, there exists $M = M(\varepsilon)$ such that $\kappa(n,e) < C\frac{e^3}{n^2}(1+\varepsilon)$, whenever $\min\{n, e/n, n^2/e\} > M$.*

Proof. Let $A > \frac{10^4}{\varepsilon^2}$ be a rational number satisfying

$$C(1-\frac{\varepsilon}{10}) < \frac{\gamma[A]}{A^3} < C(1+\frac{\varepsilon}{10}).$$

Let $M_1 = M_1(\varepsilon) \geq A$ such that, if $n > M_1$ and $e = nA$, then

$$CA^3n(1-\frac{\varepsilon}{5}) < \kappa(n,e) < CA^3n(1+\frac{\varepsilon}{5}).$$

Let $G_1 = G_1(n_1, e_1)$ be a graph with $n_1 > M_1$ vertices, $e_1 = An_1$ edges, and suppose that G_1 is drawn in the plane with $\kappa(n_1, e_1)$ crossings, where $CA^3n_1(1-\frac{\varepsilon}{5}) < \kappa(n_1,e_1) < CA^3n_1(1+\frac{\varepsilon}{5})$. For each vertex v of G_1 with degree $d(v) > A^{3/2}$, we do the following. Let $d(v) = rA^{3/2} + s$, where $0 \leq s < A^{3/2}$. Substitute v with $r+1$ vertices, each of degree $A^{3/2}$, except one which has degree s, each drawn very close to the original position of v. Clearly, this can be done without creating any additional crossing. We obtain a graph $G_2(n_2, e_2)$ such that

$$n_1 \leq n_2 \leq n_1(1+\frac{2}{\sqrt{A}}) \leq n_1(1+\frac{\varepsilon}{10}),$$

$e_2 = e_1$, and G_2 is drawn in the plane with $\kappa(n_1, e_1)$ crossings.

Suppose that n and e are fixed, $\min\{n, e/n, n^2/e\} > M(\varepsilon) = \frac{10M_1}{\varepsilon}$. Let

$$L = \frac{e/n}{e_2/n_2} \quad \text{and} \quad K = \frac{n^2/e}{n_2^2/e_2},$$

so that

$$n = KLn_2 \quad \text{and} \quad e = KL^2e_2.$$

Let

$$\tilde{L} = \left\lfloor L(1+\frac{\varepsilon}{10}) \right\rfloor \quad \text{and} \quad \tilde{K} = \left\lfloor K(1-\frac{\varepsilon}{10}) \right\rfloor$$

and let

$$\tilde{n} = \tilde{K}\tilde{L}n_2 \text{ and } \tilde{e} = \tilde{K}\tilde{L}^2 e_2.$$

Then $n(1-\frac{\varepsilon}{5}) < \tilde{n} < n$ and $e_2 < \tilde{e} \le e_2(1+\frac{\varepsilon}{4})$, so we have $\kappa(n,e) < \kappa(\tilde{n},\tilde{e})$.

Substitute each vertex of G_2 with $\tilde{L}$ very close vertices, and substitute each edge of G_2 with the corresponding $\tilde{L}^2$ edges, all running very close to the original edge. Make $\tilde{K}$ copies of this drawing, each separated from the others. This way we got a graph $\tilde{G}(\tilde{n},\tilde{e})$ drawn in the plane. We estimate the number of crossings X in this drawing.

A crossing in the original drawing of G_2 corresponds to $\tilde{K}\tilde{L}^4$ crossings in the present drawing of $\tilde{G}$. For any two edges of G_2 with common endpoint, uv and uw, the edges arise from them have at most $\tilde{K}\tilde{L}^4$ crossings with each other. So

$$X \le \tilde{K}\tilde{L}^4 \left(\kappa(n_1,e_1) + \sum_{v\in V(G_2)} \binom{d(v)}{2} \right)$$

But $\sum_{v\in V(G_2)} d(v) = 2e_2$ and $d(v) \le A^{3/2}$, so

$$\sum_{v\in V(G_2)} \binom{d(v)}{2} < 3A^{5/2}n_2.$$

Therefore,

$$\kappa(n,e) < \kappa(\tilde{n},\tilde{e}) \le c < \tilde{K}\tilde{L}^4\kappa(n_1,e_1) + \tilde{K}\tilde{L}^4 3A^{5/2}n_2 < \tilde{K}\tilde{L}^4\kappa(n_1,e_1)(1+\frac{\varepsilon}{10})$$

$$< \tilde{K}\tilde{L}^4 C A^3 n_1 (1+\frac{\varepsilon}{5})(1+\frac{\varepsilon}{10}) = \tilde{K}\tilde{L}^4 C\frac{e_1^3}{n_1^2}(1+\frac{\varepsilon}{5})(1+\frac{\varepsilon}{10})$$

$$< KL^4 C\frac{e_2^3}{n_2^2}(1+\frac{\varepsilon}{10})^6(1+\frac{\varepsilon}{5})(1+\frac{\varepsilon}{10}) < C(1+\varepsilon)\frac{e^3}{n^2}. \quad \square$$

Remark 4.4. It was shown in [PT97] that $.06 \ge C \ge .029$.

We cannot decide whether Theorem 4 remains true under the weaker condition that $C_1 n \le e \le C_2 n^2$ for suitable positive constants C_1 and C_2. If the answer were in the affirmative, then, clearly, $C_1 > 3$. We would also have that $C_2 < 1/2$, because, by [G72], for $e = \binom{n}{2}$, $\mathrm{cr}(K_n) > (\frac{1}{10} - \varepsilon)\frac{e^3}{n^2}$ for any $\varepsilon > 0$ if n is large enough.

5 Midrange crossing constants on other surfaces – Proof of Theorem 5

Lemma 5.1. *For any integer $g \ge 0$ and for any $1 > \varepsilon > 0$, there exists $N = N(g,\varepsilon)$ such that $\kappa_g(n,e) > C\frac{e^3}{n^2}(1-\varepsilon)$, whenever $\min\{n, e/n, n^{3/2}/e\} > N$.*

Proof. For $g = 0$, the assertion follows from Lemma 4.2. Suppose that $g > 0$ is fixed and we have already proved the lemma for $g-1$. For any $\varepsilon > 0$, let $N(g, \varepsilon) = \frac{10^5}{\varepsilon^2} gN(g-1, \varepsilon/10)$. Suppose, in order to get a contradiction, that $\min\{n, e/n, n^{3/2}/e\} > N$, and let $G(n, e)$ be a graph drawn on S_g with $\mathrm{cr}_g(G) = \kappa_g(n, e) < C\frac{e^3}{n^2}(1-\varepsilon)$ crossings.

As long as there is an edge with at least $4C\frac{e^2}{n^2}$ crossings, delete it. Let the resulting graph be $G_1(n_1, e_1)$. Suppose that we deleted e' edges. Then G_1 has $n_1 = n$ vertices, $e_1 = e - e'$ edges, and the number of crossings in the resulting drawing of G_1 is at most $\mathrm{cr}_g(G) - 4C\frac{e^2}{n^2}e'$. Therefore, $e' < e/4$, so $e \geq e_1 \geq 3e/4$. It is not hard to check that $\mathrm{cr}_g(G_1) < C\frac{e_1^3}{n_1^2}(1-\varepsilon)$ and G_1 contains no edge with more than $4C\frac{e^2}{n^2} < 8C\frac{e_1^2}{n_1^2}$ crossings.

Consider all cycles of G_1, as they are drawn on S_g. If each cycle is *trivial*, i.e., each cycle is contractible to a point of S_g, then every connected component of G is contractible to a point. That is, in this case, our drawing of G on S_g is equivalent to a drawing of G_1 on the plane. Consequently, $\mathrm{cr}_{g-1}(G_1) \leq \mathrm{cr}_0(G_1) < C\frac{e^3}{n^2}(1-\varepsilon)$ contradicting the induction hypothesis.

Suppose that there is a non-trivial (i.e., non-contractible) cycle $\mathcal{C}$ of G_1 with at most $\frac{\varepsilon}{80C}\frac{n_1^2}{e_1}$ edges. Clearly, $\mathcal{C}$ contains a non-trivial closed curve, $\mathcal{C}'$, which does not intersect itself. The total number of crossings along $\mathcal{C}'$ is at most

$$\frac{\varepsilon}{80C}\frac{n_1^2}{e_1}8C\frac{e_1^2}{n_1^2} = \frac{\varepsilon}{10}e_1.$$

Delete all edges that cross $\mathcal{C}'$. Cut S_g along $\mathcal{C}'$. Replace every vertex (resp. edge) $\mathcal{C}'$ by two vertices, one on each side of the cut. Every edge of G arriving at a vertex v of $\mathcal{C}'$ from a given side of the cut will be connected to the copy of v lying on the same side. Thus, we obtain a graph $G_2(n_2, e_2)$, drawn with fewer than $\mathrm{cr}_g(G_1)$ crossings. Attaching a half-sphere to each side of the cut, we obtain either a surface of genus $g-1$ or two surfaces whose genuses are smaller than g. We discuss only the former case (the calculation in the latter one is very similar). Since we doubled at most $\frac{\varepsilon}{80C}\frac{n_1^2}{e_1} = \varepsilon n_1 \frac{n_1}{e_1}\frac{1}{80C} < \varepsilon n_1 \frac{1}{N} < n_1\frac{\varepsilon}{10}$ vertices and deleted at most $\frac{\varepsilon}{10}e$ edges, we have $n_2 \leq n_1(1+\frac{\varepsilon}{10})$ and $e_2 \geq e_1(1-\frac{\varepsilon}{10})$. In the resulting drawing there are fewer than $\mathrm{cr}_g(G_1)$ crossings, therefore

$$\mathrm{cr}_{g-1}(G_2) < \mathrm{cr}_g(G_1) < C\frac{e_1^3}{n_1^2}(1-\varepsilon) \leq C\frac{e_2^3}{n_2^2}(1-\varepsilon)(1-\frac{\varepsilon}{10})^{-3}(1+\frac{\varepsilon}{10})^2 \leq C\frac{e_2^3}{n_2^2}(1-\frac{\varepsilon}{10}),$$

contradicting the induction hypothesis.

Thus, we can assume that every non-trivial cycle of G_1 contains at least $\frac{\varepsilon}{80C}\frac{n_1^2}{e_1}$ edges. For each vertex v of G_1 with degree $d(v) > \frac{10e_1}{\varepsilon n_1}$, we do the following. Let $d(v) = r\frac{10e_1}{\varepsilon n_1} + s$, where $0 \leq s < \frac{10e_1}{\varepsilon n_1}$. Without creating any new crossing, replace v by $r+1$ nearby vertices, each of degree $\frac{10e_1}{\varepsilon n}$, except one, whose degree is s. We obtain a graph $G_3(n_3, e_3)$ drawn on S_g with $n_1 \leq n_3 \leq n_1(1+\frac{\varepsilon}{5})$, $e_3 = e_1$, and with the same number of crossings as G_1. Hence,

$$\mathrm{cr}_g(G_3) \leq \mathrm{cr}_g(G_1) \leq C\frac{e_1^3}{n_1^2}(1-\varepsilon) \leq C\frac{e_3^3}{n_3^2}(1-\varepsilon)(1+\frac{\varepsilon}{5})^2 \leq C\frac{e_3^3}{n_3^2}(1-\frac{\varepsilon}{2}).$$

The maximum degree D in G_3 cannot exceed $\frac{10e_1}{\varepsilon n_1} < \frac{18e_3}{\varepsilon n_3}$, and the length of each non-trivial cycle is at least $\frac{\varepsilon}{80C}\frac{n_1^2}{e_1} \geq \frac{\varepsilon}{100C}\frac{n_3^2}{e_3}$. Apply to G_3 the DECOMPOSITION ALGORITHM described in Section 2 with the difference that, instead of (1), use the following stopping rule: STOP in STEP $i+1$ if

$$(2/3)^i < \frac{\varepsilon}{100C}\frac{n_3}{e_3}.$$

Suppose that the algorithm terminates in STEP $k+1$. Then

$$(2/3)^k < \frac{\varepsilon}{100C}\frac{n_3}{e_3} \leq (2/3)^{k-1}.$$

First, we give an upper bound on the total number of edges deleted from G_3. Let $G^0 = G_1^0 = G_3$ and $m_0 = 1$. Using (2), we obtain that, for every $0 \leq i < k$,

$$\sum_{j=1}^{m_i}\sqrt{\mathrm{cr}_g(G_j^i)} \leq \sqrt{m_i\sum_{j=1}^{m_i}\mathrm{cr}_g(G_j^i)} \leq \sqrt{(3/2)^{i+1}}\sqrt{\mathrm{cr}_g(G_3)} \leq \sqrt{(3/2)^{i+1}}\sqrt{C\frac{e_3^3}{n_3^2}(1-\frac{\varepsilon}{2})}.$$

Denoting by $d(v, G_j^i)$ the degree of vertex v in G_j^i, we have

$$\sum_{j=1}^{m_i}\sqrt{\sum_{v\in V(G_j^i)} d^2(v,G_j^i)} \leq \sqrt{(3/2)^{i+1}}\sqrt{\sum_{v\in V(G^i)} d^2(v,G^i)}$$

$$\leq \sqrt{(3/2)^{i+1}}\sqrt{\max_{v\in V(G^i)} d(v,G^i)\sum_{v\in V(G^i)} d(v,G^i)} \leq \sqrt{(3/2)^{i+1}}\sqrt{\frac{18e_3^3}{\varepsilon n_3^2}(2e_3)} = 12\sqrt{(3/2)^{i+1}}\frac{e_3}{\sqrt{\varepsilon n_3}}.$$

By Theorem 6 (proved in the last section), the total number of edges deleted during the algorithm is

$$\sum_{i=0}^{k-1}\sum_{j=1}^{m_i} b(G_j^i) \leq 300(1+g^{3/4})\sum_{i=0}^{k-1}\sum_{j=1}^{m_i}\sqrt{\mathrm{cr}_g(G_j^i) + \sum_{v\in V(G_j^i)} d^2(v,G_j^i)}$$

$$\leq 300(1+g^{3/4})\sum_{i=0}^{k-1}\sum_{j=1}^{m_i}\sqrt{\mathrm{cr}_g(G_j^i)} + 300(1+g^{3/4})\sum_{i=0}^{k-1}\sum_{j=1}^{m_i}\sqrt{\sum_{v\in V(G_j^i)} d^2(v,G_j^i)}$$

$$\leq 300(1+g^{3/4})\sum_{i=0}^{k-1}\sqrt{(3/2)^{i+1}}\left(\sqrt{C\frac{e_3^3}{n_3^2}(1-\frac{\varepsilon}{2})} + 6\frac{e_3}{\sqrt{\varepsilon n_3}}\right)$$

$$\leq 300(1+g^{3/4})\sqrt{3/2}\frac{\sqrt{(3/2)^k}-1}{\sqrt{3/2}-1}\left(\sqrt{C\frac{e_3^3}{n_3^2}(1-\frac{\varepsilon}{2})} + 6\frac{e_3}{\sqrt{\varepsilon n_3}}\right)$$

$$\leq 2000(1+g^{3/4})\sqrt{\frac{C}{\varepsilon}}\sqrt{\frac{e}{n}}\left(\sqrt{C\frac{e_3^3}{n_3^2}(1-\frac{\varepsilon}{2})} + 6\frac{e_3}{\sqrt{\varepsilon n_3}}\right) \leq e_3\frac{\varepsilon}{10}.$$

Therefore, the number of edges $e(G^k)$ of the graph G^k obtained in the final STEP of the algorithm satisfies $e(G^k) \geq e_3(1-\frac{\varepsilon}{10})$. Consider the drawing of G^k on S_g inherited from the drawing of G_3. Each connected component of G^k has fewer than $\frac{\varepsilon}{100C}\frac{n_3^2}{e_3}$ vertices, therefore, each cycle of G^k, as drawn on S_g, is contractible to a point. Consequently, this drawing is equivalent to a planar drawing of G^k. Hence,

$$\mathrm{cr}_{g-1}(G^k) \leq \mathrm{cr}_0(G^k) \leq \mathrm{cr}_g(G_3) \leq C\frac{e_3^3}{n_3^2}(1-\frac{\varepsilon}{2}) \leq C\frac{e^3(G^k)}{n^2(G^k)}(1-\frac{\varepsilon}{2})(1-\frac{\varepsilon}{10})^{-3}$$

$$< C\frac{e^3(G^k)}{n^2(G^k)}(1-\frac{\varepsilon}{10}),$$

a contradiction. This concludes the proof of Lemma 5.1. □

Lemma 5.2. *For any integer $g \geq 0$ and for any $\varepsilon > 0$, there exists $N' = N'(g,\varepsilon)$ such that $\kappa_g(n,e) > C\frac{e^3}{n^2}(1-\varepsilon)$, whenever $\min\{n, e/n, n^2/e\} > N'$.*

Proof. The proof is analogous to that of Lemma 4.2. □

Lemma 5.3. *For any integer $g \geq 0$ and for any $\varepsilon > 0$, there exists $M = M(g,\varepsilon)$ such that $\kappa_g(n,e) < C\frac{e^3}{n^2}(1+\varepsilon)$, whenever $\min\{n, e/n, n^2/e\} > M$.*

Proof. Clearly, for any graph G and for any $g \geq 0$, we have $\mathrm{cr}_0(G) \geq \mathrm{cr}_g(G)$. Therefore, Lemma 5.3 is a direct consequence of Lemma 4.3. □

Theorem 5 now readily follows from Lemmas 5.2 and 5.3.

6 A separator theorem – Proof of Theorem 6

For the proof of Theorem 6, we need a slight variation of the notion of bisection width. The *weak bisection width*, $\overline{b}(G)$, of a graph G is defined as the minimum number of edges whose removal splits the graph into two components, each of size at least $|V(G)|/5$. That is,

$$\overline{b}(G) = \min_{|V_A|,|V_B| \geq n/5} |E(V_A, V_B)|,$$

where $E(V_A, V_B)$ denotes the number of edges between V_A and V_B, and the minimum is taken over all partitions $V(G) = V_A \cup V_B$ with $|V_A|, |V_B| \geq |V(G)|/5$.

Lemma 6.1 *For any graph G, we have*

$$\overline{b}(G) \leq b(G) \leq 2\max_{H \subset G} \overline{b}(H).$$

Proof. The first inequality is obviously true. To prove the second one, let $|V(G)| = n$ and consider a partition $V(G) = V_A \cup V_B$ such that $n/5 \leq |V_A|, |V_B| \leq 4n/5$ and $|E(V_A, V_B)| =$

$\overline{b}(G)$. Suppose that $|V_A| \le |V_B|$. If $n/3 \le |V_A|$, then $b(G) = \overline{b}(G)$ and we are done. So we can assume that $n/5 \le |V_A| \le n/3$ and $2n/3 \le |V_B| \le 4n/5$.

Let H be the subgraph of G induced by V_B. By definition, there is a partition $V_B = V_B' \cup V_B''$ such that $|V_B|/5 \le |V_B'|, |V_B''| \le 4|V_B|/5$ and $|E(V_B', V_B'')| = \overline{b}(H)$. We can assume that $|V_B'| \le |V_B''|$. Then

$$\frac{n}{3} \le \frac{|V_B|}{2} \le |V_B''| \le \frac{4|V_B|}{5} \le \frac{16n}{25} < \frac{2n}{3}.$$

Letting $V_1 = V_A \cup V_B'$ and $V_2 = V_B''$, we have $V(G) = V_1 \cup V_2$, $n/3 \le |V_1|, |V_2| \le 2n/3$,

$$|E(V_1, V_2)| \le |E(V_A, V_B)| + |E(V_B', V_B'')| \le \overline{b}(G) + \overline{b}(H),$$

and the result follows. □

Theorem 6 is an immediate consequence of Lemma 6.1 and the following statement.

Theorem 6.2. *Let G be a graph with n vertices of degrees $d_1, d_2, \ldots, d_n$. Then*

$$\overline{b}(G) \le 150(1 + g^{3/4})\sqrt{cr_g(G) + \sum_{i=1}^{n} d_i^2}.$$

Proof. Clearly, we can assume that G contains no isolated vertices, that is, $d_i > 0$ for all $1 \le i \le n$. Consider a drawing of G on S_g with exactly $\mathrm{cr}_g(G)$ crossings. Let $v_1, v_2, \ldots, v_n$ be the vertices of G with degrees $d_1, d_2, \ldots, d_n$, respectively. Introduce a new vertex at each crossing. Denote the set of these vertices by V_0. Replace each $v_i \in V(G)$ $(i = 1, 2 \ldots, n)$ by a set V_i of vertices forming a $d_i \times d_i$ piece of a square grid, in which each vertex is connected to its horizontal and vertical neighbors. Let each edge incident to v_i be hooked up to distinct vertices along one side of the boundary of V_i without creating any crossing. These d_i vertices will be called the *special boundary vertices* of V_i.



Figure 1.

Thus, we obtain a graph H of $\sum_{i=0}^{n} |V_i| = \mathrm{cr}_g(G) + \sum_{i=1}^{n} d_i^2$ vertices and no crossing (see Fig. 1.). For each $1 \leq i \leq n$, assign weight $1/d_i$ to each special boundary vertex of V_i. Assign weight 0 to all other vertices of H. For any subset ν of the vertex set of H, let $w(\nu)$ denote the total weight of the vertices belonging to ν. With this notation, $w(V_i) = 1$ for each $1 \leq i \leq n$. Consequently, $w(V(H)) = n$.

Since H is drawn on S_g without crossing, H does not contain K_α as a minor, where $\alpha = \lfloor 4 + 4\sqrt{g} \rfloor$ [RY68]. Then, by a result of Alon, Seymour, and Thomas [AST90], the vertices of H can be partitioned into three sets, A, B and C, such that $w(A), w(B) \geq n/3$ and $|C| \leq 25(1+g^{3/4})\sqrt{\mathrm{cr}_g(G) + \sum_{i=1}^{n} d_i^2}$, and there is no edge from A to B. Let $A_i = A \cap V_i$, $B_i = B \cap V_i$, $C_i = C \cap V_i$ $(i = 0, 1, \ldots, n)$.

For any $1 \leq i \leq n$, we say that V_i is of *type A* (resp. *type B*) if $w(A_i) \geq 5/6$ (resp. $w(B_i) \geq 5/6$), and it is of *type C,* otherwise.

Define a partition $V(G) = V_A \cup V_B$ of the vertex set of G, as follows. For any $1 \leq i \leq n$, let $v_i \in V_A$ (resp. $v_i \in V_B$) if V_i is of type A (resp. type B). The remaining vertices, $\{v_i \mid V_i \text{ is of type } C\}$ are assigned either to V_A or to V_B so as to minimize $||V_A| - |V_B||$.

Claim 1. $n/5 \leq |V_A|, |V_B| \leq 4n/5$

To prove the claim, define another partition $V(H) = \overline{A} \cup \overline{B} \cup \overline{C}$ such that $\overline{A} \cap V_i = A \cap V_i$ and $\overline{B} \cap V_i = B \cap V_i$, for $i = 0$ and for every V_i of type C. If V_i is of type A (resp. type B), then let $V_i = \overline{A}_i \subset \overline{A}$ (resp. $V_i = \overline{B}_i \subset \overline{B}$), finally, let $\overline{C} = V(H) - \overline{A} - \overline{B}$.

For any V_i of type A, $w(\overline{A}_i) - w(A_i) \leq w(A_i)/5$. Similarly, for any V_i of type B, $w(\overline{B}_i) - w(B_i) \leq w(B_i)/5$. Therefore,

$$|w(\overline{A}) - w(A)| \leq \max\{w(A), w(B)\}/5 \leq 2n/15.$$

Hence, $n/5 \leq w(\overline{A}) \leq 4n/5$ and, analogously, $n/5 \leq w(\overline{B}) \leq 4n/5$. In particular, $|w(\overline{A}) - w(\overline{B})| \leq 3n/5$. Using the minimality of $||V_A| - |V_B||$, we obtain that $||V_A| - |V_B|| \leq 3n/5$, which implies Claim 1.



Figure 2.

Claim 2. *For any* $1 \leq i \leq n$,
(i) *if* V_i *is of type A (resp. of type B), then* $w(B_i)d_i \leq |C_i|$ *(resp.* $w(A_i)d_i \leq |C_i|$*);*
(ii) *if* V_i *is of type C, then* $d_i/6 \leq |C_i|$.

In V_i, every connected component belonging to A_i is separated from every connected component belonging to B_i by vertices in C_i. There are $w(A_i)d_i$ (resp. $w(B_i)d_i$) special boundary vertices in V_i, which belong to A_i (resp. B_i). It can be shown by an easy case analysis that the number of separating points $|C_i| \geq \min\{w(A_i), w(B_i)\}d_i$, and Claim 2 follows (see Fig. 2.).

In order to establish Theorem 6.2 (and hence Theorem 6), it remains to prove the following statement.

Claim 3. *The total number of edges between* V_A *to* V_B *satisfies*

$$|E(V_A, V_B)| \leq 150(1 + g^{3/4})\sqrt{\mathrm{cr}_g(G) + \sum_{i=1}^{n} d_i^2}.$$

To see this, denote by E_0 the set of all edges of H adjacent to at least one element of C_0. For any $1 \leq i \leq n$, define $E_i \subset E(H)$ as follows. If V_i is of type A (resp. type B), let E_i consist of all edges leaving V_i and adjacent to a special boundary vertex belonging to B_i (resp. A_i). If V_i is of type C, let all edges leaving V_i belong to E_i.

For any $1 \leq i \leq n$, let E_i' denote the set of edges of G corresponding to the elements of E_i $(0 \leq i \leq n)$. Clearly, we have $|E_i'| \leq |E_i|$, because distinct edges of G give rise to distinct edges of H. It is easy to see that every edge between V_A and V_B belongs to $\cup_{i=0}^{n} E_i'$.

Obviously, $|E_0'| \leq |E_0| \leq 4|C_0|$. By Claim 2, if V_i is of type A or of type B, then $|E_i'| \leq |E_i| \leq |C_i|$. If V_i is of type C, then $|E_i'| \leq |E_i| = d_i \leq 6|C_i|$. Therefore,

$$|E(V_A, V_B)| \leq |\cup_{i=0}^{n} E_i'| \leq \sum_{i=0}^{n} |E_i| \leq 6|C| \leq 150(1 + g^{3/4})\sqrt{\mathrm{cr}_g(G) + \sum_{i=1}^{n} d_i^2}.$$

This concludes the proof of Claim 3 and hence Theorem 6.2 and Theorem 6. □

Acknowledgement. We would like to express our gratitude to Zoltán Szabó for his help in writing Section 5, and to László Székely for many very useful remarks.

References

[ACNS82] M. Ajtai, V. Chvátal, M. Newborn, and E. Szemerédi, Crossing-free subgraphs, in: *Theory and Practice of Combinatorics, North-Holland Math. Stud.* **60,** North-Holland, Amsterdam-New York, 1982, 9–12.

[ARS98] N. Alon, L. Rónyai, and T. Szabó, Norm-graphs: variations and applications, to appear.

[AST90] N. Alon, P. Seymour, and R. Thomas, A separator theorem for nonplanar graphs, *J. Amer. Math. Soc.* **3** (1990), 801–808.

[AST94] N. Alon, P. Seymour, and R. Thomas, Planar separators, *SIAM J. of Discrete Math.* **7** (1994), 184–193.

[Be66] C. Benson, Minimal regular graphs of girth eight and twelve, *Canadian J. Mathematics* **18** (1966), 1091–1094.

[BS74] J. A. Bondy and M. Simonovits, Cycles of even length in graphs, *J. Combinatorial Theory, Ser. B* **16** (1974), 97–105.

[B66] W. G. Brown, On graphs that do not contain a Thomsen graph, *Canadian Mathematical Bulletin* **9** (1966), 281-285.

[D98] T. K. Dey, Improved bounds for planar k-sets and related problems, *Discrete and Computational Geometry* **19** (1998), 373–382.

[EG73] P. Erdős and R. K. Guy, Crossing number problems, *American Mathematical Monthly* **80** (1973), 52–58.

[ER62] P. Erdős and A. Rényi, On a problem in the theory of graphs, (in Hungarian), *Magyar Tud. Akad. Mat. Kutató Int. Közl.* **7** (1962), 623–641.

[F96] Z. Füredi, New asymptotics for bipartite Turán numbers, *J. Combinatorial Theory, Ser. A* **75** (1996), 141–144.

[GJ83] M. R. Garey and D. S. Johnson, Crossing number is NP-complete, *SIAM J. Alg. Disc. Meth.* **4** (1983), 312–316.

[G72] R. K. Guy, Crossing numbers of graphs, in: *Graph theory and applications (Proc. Conf. Western Michigan Univ., Kalamazoo, Mich., 1972)* Lecture Notes in Mathematics **303,** Springer, Berlin, 111-124.

[KST54] T. Kővári, V. T. Sós, and P. Turán, On a problem of K. Zarankiewicz, *Colloquium Mathematicum* **3** (1954), 50–57.

[L83] F. T. Leighton, *Complexity Issues in VLSI,* MIT Press, Cambridge, 1983.

[L84] F. T. Leighton, New lower bound techniques for VLSI, *Math. Systems Theory* **17** (1984), 47–70.

[LT79] R. Lipton and R. Tarjan, A separator theorem for planar graphs, *SIAM J. Applied Mathematics* **36** (1979), 177–189.

[PSS96] J. Pach, F. Shahrokhi, and M. Szegedy, Applications of the crossing number, *Algorithmica* **16** (1996), 111–117.

[PS98] J. Pach and M. Sharir, On the number of incidences between points and curves, *Combinatorics, Probability and Computing* **7** (1998), 121–127.

[PT97] J. Pach and G. Tóth, Graphs drawn with few crossings per edge, *Combinatorica* **17** (1997), 427–439.

[R58] I. Reiman, Über ein Problem von K. Zarankiewicz, *Acta Math. Acad. Sci. Hungar.* **9** (1958), 269–279.

[RT97] R. B. Richter and C. Thomassen, Relations between crossing numbers of complete and complete bipartite graphs, *American Mathematical Monthly*, February 1997, 131–137.

[RY68] G. Ringel and J. W. T. Youngs, Solution of the Heawood map-coloring problem, *Proc. Nat. Acad. Sci. U.S.A.* **60** (1968), 438–445.

[S97] M. Simonovits, personal communication.

[S66] R. Singleton, On minimal graphs of maximum even girth, *J. Combinatorial Theory* **1** (1966), 306–332.

[S98] L. A. Székely, Crossing numbers and hard Erdős problems in discrete geometry, *Combinatorics, Probability and Computing* **6** (1998), 353–358.

[W91] R. Wenger, Extremal graphs with no C_4's, C_6's, or C_{10}'s, *J. Combinatorial Theory, Ser. B* **52** (1991), 113–116.

[WB78] A. T. White and L. W. Beineke, Topological graph theory, in: *Selected Topics in Graph Theory (L. W. Beineke and R. J. Wilson., eds.),* Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], London-New York, 1983, 15–49.