DIMACS Technical Report 99-46
October 1999

Universal $H-$colorable Graphs Without A Given Configuration

by

Paul A. Dreyer, Jr.[1]
Dept. of Computer Science
Rutgers University
New Brunswick, New Jersey 08903

Christopher D. Malon[2]
Dept. of Mathematics
University of Chicago
Chicago, IL 60637

Jaroslav Nešetřil[3]
Dept. of Applied Mathematics
Charles University
Prague, Czech Republic

[1]Affiliated Graduate Student Member
[2]DIMACS REU Participant
[3]Permanent DIMACS Member

DIMACS is a partnership of Rutgers University, Princeton University, AT&T Labs-Research, Bell Labs, Telcordia Technologies (formerly Bellcore) and NEC Research Institute.

DIMACS is an NSF Science and Technology Center, funded under contract STC–91–19999; and also receives support from the New Jersey Commission on Science and Technology.

ABSTRACT

For every pair of finite connected graphs F and H, and every integer k, we construct a universal graph U with the following properties:

1. There is a homomorphism $\pi : U \to H$, but no homomorphism from F to U.
2. For every graph G with maximal degree no more than k having a homomorphism $h : G \to H$, but no homomorphism from F to G, there is a homomorphism $\alpha : G \to U$, such that $h = \pi \circ \alpha$.

Particularly, this solves a problem presented in [1] and [2] regarding the chromatic number of a universal graph.

1 Introduction

For a graph G, let $V(G)$ and $E(G)$ denote its vertex and edge sets, respectively. Given two graphs G and H, a map $f : V(G) \to V(H)$ is a *(graph) homomorphism* from G to H if and only if $\{f(x), f(y)\} \in E(H)$ for all edges $\{x, y\} \in E(G)$. For simplicity, $f : G \to H$ will denote that f is a homomorphism from G to H, and $G \not\to H$ means that no homomorphism exists from G to H.

Homomorphisms extend the notion of coloring, as $\chi(G) \leq k$ if and only if $G \to K_k$. More generally, we say that G is *H-colorable* if there is a homomorphism $G \to H$. If $F \not\to G$, we say that G is *F-free*. A graph U is *universal* for a class of graphs $\mathcal{K}$ if, for any graph $G \in \mathcal{K}$, there exists a homomorphism $G \to U$. For example, the complete graph on k vertices is universal for the family of k-colorable graphs. Universal graphs were considered earlier in [1] and [2], which established their existence for several classes of graphs, including classes of $F-$free graphs of bounded degree.

When considering $F-$free and $H-$colorable graphs, we can restrict to those graphs F and H that are *cores*, graphs for which every endomorphism is an automorphism. Indeed, a graph is $F-$free if and only if it is free of the (uniquely determined) core of F, and $H-$colorable if and only if it is colorable by the core of H, [3]. We have the following theorem from [5]:

Theorem 1.1 *Let F, H be fixed cores. Then for every universal graph U of the class of all $F-$free, $H-$colorable graphs K_H, there exists a homomorphism $H \to U$.*

In particular, every universal graph for 3-colorable, triangle-free graphs contains a triangle.

This negative result was complemented by Häggkvist and Hell with a positive result for $F-$free graphs of bounded degree [2]. We say that a graph is *$b-$bounded* if $\Delta(G) \leq b$.

Theorem 1.2 *Let F be a core, b a positive integer. Then the class $\mathcal{K}_F^b$ of all $b-$bounded $F-$free graphs has a universal $F-$free graph U_F^b.*

As the chromatic number of a $b-$bounded graph is $\leq b+1$, all of the graphs in the class $\mathcal{K}_F^b$ have chromatic number $\leq b+1$. Häggkvist and Hell [2] have asked the minimal chromatic number of an universal graph for K_F^b. The graphs constructed in [2] all have a large chromatic number. In [1], Galluccio, Hell, and Nešetřil considered the complexity of $H-$coloring 3-bounded graphs with $\chi(H) \leq 3$, and conjectured that $H-$coloring of 3-bounded graphs is an NP-complete problem for any 3-chromatic triangle-free graph H.

In this paper, we answer both of these questions by proving that the class of all $F-$free 3-bounded graphs has a universal $F-$free graph U with $\chi(U) \leq 3$. More generally, our main result, Theorem 3.1, demonstrates the existence of $F-$free, $H-$colorable universal graphs for the family of $b-$bounded, $F-$free, $H-$colorable graphs.

By a result of Johanson [4], we have the following corollary:

Corollary 1.3 *For every positive integer b, there exists a graph U_b with the following properties:*

1. *U_b is triangle-free, and $\chi(U_b) \leq c \cdot \frac{b}{\log b}$ for a constant c independent of b.*

2. *U_b is universal for the family of $b-$bounded, triangle-free graphs.*

The paper is organized as follows. In Section 2, we give a proof of a theorem covering a particular case of the main theorem (when $F = K_3 = H$). This construction closely follows the construction given in [1] and [2]. In Section 3, we prove the general theorem, giving a new construction for the above theorem. Section 4 contains some remarks and open problems.

We introduce some notation used for the remainder of the paper. As usual, $[n]$ will denote the set of positive integers $\{1, 2, \ldots, n\}$. If x and y are two vertices in a connected graph G, the *distance* $d_G(x, y)$ denotes the number of edges in the shortest path in G joining x and y. For a graph G and a non-negative integer k, G^k is a graph defined by $V(G^k) = V(G)$ and $E(G^k) = \{\{x, y\} : x, y \in V(G), 1 \leq d_G(x, y) \leq k\}$. For a vertex v, $G[v, k]$ denotes the subgraph induced by the vertices $w \in V(G)$ such that $0 \leq d_G(v, w) \leq k$.

If $\Delta(G) = k$, then by induction, for a vertex $v \in G$ and $j \geq 1$, at most $k(k-1)^{j-1}$ points have distance j from v. Putting $X_k(d) = 1 + \sum_{j=1}^{d} k(k-1)^{j-1}$, it follows that $|V(G[v, d])| \leq X_k(d)$.

2 $k-$colorable, triangle-free universal graphs for $k-$bounded, triangle-free graphs

Theorem 2.1 *Given a positive integer k, let $\mathcal{F}$ be the family of $k-$bounded, triangle-free graphs. There exists a universal graph U such that:*

1. *$\chi(U) \leq k$.*

2. *For every $G \in \mathcal{F}$, there exists a homomorphism $\alpha : G \to U$.*

3. *U is triangle-free.*

Proof: Let $G \in \mathcal{F}$. Assume without loss of generality that G is k-regular. Since $\chi(G) \leq k$, we know there exists a proper coloring of G, $c_1 : V(G) \to [k]$. For any vertex in G^3, the degree is at most $X_k(3) - 1$, so there exists a proper coloring of G^3, $c_2 : V(G^3) = V(G) \to [X_k(3)]$. Putting $X = [k] \times [X_k(3)]$, we can define a map $c : V(G) \to X$ by $c(x) = (c_1(x), c_2(x))$. Let $\pi_1 : X \to [k]$ be the projection onto the first coordinate $\pi_1(c_1, c_2) = c_1$.

The universal graph U is defined by:

$$V(U) = X \times \binom{X}{k}$$

$$\begin{aligned} E(U) = \{\{(x, A), (x', A')\} : & A \cap A' = \emptyset, \\ & x \in A', x' \in A, \\ & \pi_1(x) \neq \pi_1(x')\}. \end{aligned}$$

1. $\chi(U) \leq k$.

   The map $\pi : V(U) \to [k]$ given by $\pi((x, A)) = \pi_1(x)$ $k-$colors U.

2. For every $G \in \mathcal{F}$, there exists a homomorphism $\alpha : G \to U$.

   Put $\alpha(v) = (c(v), \{c(w) : \{v, w\} \in E(G)\})$. If $\{v, w\} \in E(G)$ with $\alpha(v) = (x, A)$, $\alpha(w) = (x', A')$, then:

   a. $A \cap A' = \emptyset$: Because G is triangle-free, v and w have no common neighbors. No neighbor of v can receive the same $X_k(3)-$color as a neighbor of w, because neighbors of v and w are at most distance three from one another. Thus, any members of A and A' differ in their x_2-coordinate.

   b. $x \in A'$, $x' \in A$: Because $\{v, w\} \in E(G)$, $x' = c(w) \in A$ and $x = c(v) \in A'$.

   c. $\pi_1(x) \neq \pi_1(x')$: $\pi_1(x) = c_1(v)$ and $\pi_1(x') = c_1(w)$, and c_1 properly $k-$colors G, hence $\pi_1(x) \neq \pi_1(x')$.

3. U is triangle-free.

   Suppose that (x, A), (x', A'), and (x'', A'') are three vertices of U that form a triangle. Then $x \in A'$ and $x \in A''$, contradicting $A' \cap A'' = \emptyset$.

Although [1] conjectured that for triangle-free graphs H with chromatic number 3, the $H-$coloring problem for 3$-$bounded graphs is NP-complete, setting $k = 3$ in the above theorem proves this conjecture false: All 3$-$bounded, triangle-free graphs are $U-$colorable, and all 3$-$bounded graphs containing a triangle are not.

3 $F-$free universal graphs for $H-$colorable, $F-$free graphs

The preceding construction mapped triangle-free, $k-$bounded graphs into a universal graph by describing a vertex's neighborhood with the colors used by a coloring of G^3. The condition $A \cap A' = \emptyset$ sufficed to avoid triangles in the universal graph. In order to avoid more complicated graphs, we replace the coloring of G^3 with a labelled copy of the induced subgraph formed by a large neighborhood around the vertex, which will be $F-$free. Edges in the universal graph connect only the neighborhoods that might be had by adjacent vertices in $F-$free graphs. This idea allows us to strengthen the preceding theorem:

Theorem 3.1 *Let F, H be finite connected graphs, b a positive integer. Then there exists a universal graph U with the following properties:*

1. *U is $H-$colorable.*

2. *For every $b-$bounded, $F-$free, $H-$colorable graph G, there exists a homomorphism $G \to U$ that preserves the $H-$coloring of G.*

3. *U is $F-$free.*



This result can be expressed using a commutative diagram:

Proof: Throughout this proof, we assume that given a K_n, its vertices are labelled by $[n]$. Let $m = |V(F)|$, and let $\mathcal{S}$ be the set of all connected subgraphs S of $K_{X_k(m)}$ such that $F \not\to S$.

We will now describe the universal graph U.

$$\begin{aligned} V(U) &= \mathcal{S} \times V(K_{X_k(m)}) \times V(H) \\ E(U) = \{\{(S,v,n),(S',v',n')\} :& \{n,n'\} \in E(H), \\ & \{v,v'\} \in E(S) \cap E(S'), \\ & S[v,m-1] = S'[v,m-1], \\ & S[v',m-1] = S'[v',m-1]\} \end{aligned}$$

Here, graph equalities mean not only that the two graphs are isomorphic, but that the labels on the vertices and edges in each graph are identical.

1. U is $H-$colorable.

   The map $\pi : U \to H$ defined by $\pi((S,v,n)) = n$ is a homomorphism that preserves the $H-$coloring of G.

2. For every $b-$bounded, $F-$free, $H-$colorable graph G, there exists a homomorphism $G \to U$ that preserves the $H-$coloring of G.

   Let G be a graph that satisfies the conditions above. Because G^m is $X_k(m)-$colorable, we can fix a homomorphism $c : G \to K_{X_k(m)}$. If $v \in V(G)$ and $r \leq m$, then c is injective on $G[v,r]$, so $c(G[v,r])$ is isomorphic to $G[v,r]$. In particular, $F \not\to c(G[v,r])$, since $c(G[v,r])$ is isomorphic to a subgraph of G.

   Because $F \not\to c(G[v,m])$, we can define $\alpha : V(G) \to V(U)$ by $\alpha(x) = (c(G[v,m]), c(x), h(x))$. The map α preserves the $H-$coloring of G, namely $h = \pi \circ \alpha$. We claim that α

is a homomorphism from G to U. Suppose $\{x, x'\} \in E(G)$. Because h is a homomorphism, $\{h(x), h(x')\} \in E(H)$. Since $G[x, m]$ and $G[x', m]$ are induced subgraphs of G, both containing x and x', and c is a homomorphism, $\{c(x), c(x')\} \in E(c(G[x, m])) \cap E(c(G[x', m]))$. From $G[x, m-1] \subseteq G[x', m]$ and $G[x', m-1] \subseteq G[x, m]$, we obtain:

$$\begin{aligned} c(G[x,m])[c(x), m-1] &= c(G[x, m-1]) \\ &= c((G[x', m])[x, m-1]) \\ &= c(G[x', m])[c(x), m-1]. \end{aligned}$$

Similarly, $c(G[x', m])[c(x'), m-1] = c(G[x, m])[c(x'), m-1]$. Hence, $\{\alpha(x), \alpha(x')\} \in E(U)$.

3. U is $F-$free.

   Let $V(F) = \{x_1, \ldots, x_m\}$. Suppose that a homomorphism $\beta : F \to U$ exists, and $\beta(x_i) = (S_i, v_i, n_i)$. We claim that the map $\beta^* : F \to S_1$ defined by $\beta^*(x_i) = v_i$ is a homomorphism, contradicting the condition on $V(U)$ that $F \not\to S$ for all $S \in \mathcal{S}$.

   If $\{x_i, x_j\} \in E(F)$, $d_F(x_1, x_i) \leq m - 1$ because $|V(F)| = m$. Since $\{\beta(x_i), \beta(x_j)\} \in E(U)$, $\{v_i, v_j\} \in E(S_i) \cap E(S_j)$.

   If $\{x_{k_1} = x_1, x_{k_2}, \ldots, x_{k_p} = x_i\}$ is the shortest path in F from x_1 to x_i, then $\{v_i, v_j\} \in E(S_{k_p}[v_{k_p}, 1])$. Since $\{v_{k_{p-1}}, v_{k_p}\} \in E(S_{k_p}[v_{k_p}, 1])$,

$$\{v_i, v_j\} \in E(S_{k_p}[v_{k_{p-1}}, 2]) = E(S_{k_{p-1}}[v_{k_{p-1}}, 2]).$$

   By induction, we can show that for any q, $1 \leq q \leq p < m$, that

$$\{v_i, v_j\} \in E(S_{k_{p-(q-1)}}[v_{k_{p-(q-1)}}, q]).$$

   Taking $q = p$,

$$\{v_i, v_j\} \in E(S_{k_1}[v_{k_1}, p]) = E(S_1[v_1, p]) \subseteq E(S_1).$$

   Thus, β^* is a homomorphism from F to S_1, producing a contradiction and completing the proof.

4 Remarks and Open Problems

The theorems above suggest that one needs some restrictive condition to guarantee the existence of an universal graph. In the theorems above, the $b-$boundedness condition in the graph family is essential. The following problems appear to be open.

- Does there exist a triangle-free graph U which is universal for every triangle-free planar graph?
- Does there exist a triangle-free planar graph U which is universal for all 3-bounded, triangle-free planar graphs?

We can observe that there is no triangle-free planar graph U which is universal for all triangle-free planar graphs. To see this, consider "sunflower graphs" Sun_n. To construct Sun_n, begin with a $K_{1,n}$, where the lone vertex is labelled x and the n vertices are labelled by $[n]$. Vertices i and $i+1$ are joined by edge-disjoint paths of length 3, as well as vertices n and 1. Then, the neighbors of each vertex in $[n]$ (not including x) are joined by edge-disjoint paths of length three. The sunflower graph Sun_n has $5n+1$ vertices and $7n$ edges. The graphs of Sun_n are shown below for $n = 1, 2, 3$, and 6.

One can check that there is a homomorphism $Sun_n \to Sun_{n'}$ if and only if $n'|n$. One can also check that any planar homomorphic image of Sun_n must have a vertex of degree n', where $n'|n$ and $n' > 1$. If n is prime and exceeds $\Delta(U)$, then $Sun_n \not\to U$.



Figure 1: Sunflower graphs for $n = 1, 2, 3$, and 6. Note that although Sun_1 can be constructed, it is not triangle-free. Arrows denote that homomorphisms exist from one graph to the other; slashed arrows indicate that no homomorphism exists.

Acknowledgements

The authors would like to thank the NSF (Grant No. INT 98-17956), and KONTAKT (Grant No. 202-16/209019), for funding the DIMACS/DIMATIA Research Experience for Undergraduates, which brought two of the authors (Paul Dreyer, the coordinator of the DDREU, and Christopher Malon, an undergraduate at the University of Chicago) to Charles University for a three-week research program. The work of Paul Dreyer was also supported by NSF Grant No. SBR 97-09134 to Rutgers University.

References

[1] A. Galluccio, P. Hell, and J. Nešetřil. The complexity of $H-$coloring of bounded degree graphs. *Discrete Math.*, (to appear).

[2] R. Häggkvist and P. Hell. Universality of A-mote graphs. *Europ. J. Combinatorics*, 14:23–27, 1993.

[3] P. Hell and J. Nešetřil. The core of a graph. *Discrete Math.*, 109:117–126, 1992.

[4] A. Johansson. *Some Results on Colourings of Graphs.* PhD thesis, University of Umea, 1994.

[5] J. Nešetřil and V. Rödl. Chromatically optimal rigid graphs. *J. Comb. Th. B.*, 46:133–141, 1989.