DIMACS Technical Report 2000-11
March 2000

A Note on Set Systems with no Union of Cardinality 0 Modulo m

by

Vince Grolmusz[1]
Department of Computer Science
Eötvös University, H-1053 Budapest
HUNGARY
E-mail: grolmusz@cs.elte.hu

[1]Special Year Visitor at DIMACS Center, Piscataway, NJ.

DIMACS is a partnership of Rutgers University, Princeton University, AT&T Labs-Research, Bell Labs, NEC Research Institute and Telcordia Technologies (formerly Bellcore).

DIMACS is an NSF Science and Technology Center, funded under contract STC–91–19999; and also receives support from the New Jersey Commission on Science and Technology.

ABSTRACT

Alon, Kleitman, Lipton, Meshulam, Rabin and *Spencer* (Graphs. Combin. 7 (1991), no. 2, 97-99) proved, that for any hypergraph $\mathcal{F} = \{F_1, F_2, \ldots, F_{d(q-1)+1}\}$, where q is a prime-power, and d denotes the maximal degree of the hypergraph, there exists an $\mathcal{F}_0 \subset \mathcal{F}$, such that $|\bigcup_{F\in\mathcal{F}_0} F| \equiv 0 \pmod{q}$. We give a direct, alternative proof for this theorem, and we also give an explicit construction of a hypergraph of degree d and size $\Omega(d^2)$ which does not contain a non-empty sub-hypergraph with a union of size 0 modulo 6.

Keywords: composite modulus, hypergraphs, polynomials over rings

1 Introduction

Alon, Kleitman, Lipton, Meshulam, Rabin and *Spencer* [1] gave the following definition:

Definition 1 ([1]) *For integers $d, m \geq 1$, let $f_d(m)$ denote the smallest t such that for any hypergraph $\mathcal{F} = \{F_1, F_2, \ldots, F_t\}$ with maximum degree d there exists a non-empty $\mathcal{F}_0 \subset \mathcal{F}$, such that $|\bigcup_{F \in \mathcal{F}_0} F| \equiv 0 \pmod{m}$*

Baker and *Schmidt* [2] defined the following quantity:

Definition 2 *For integers $d, m \geq 1$, let $g_d(m)$ denote the smallest t such that for any polynomial $h \in Z[x_1, x_2, \ldots, x_t]$ of degree d, satisfying $h(0)=0$, there exists an $0 \neq \varepsilon \in \{0,1\}^n$, such that $h(\varepsilon) \equiv 0 \pmod{m}$.*

The following theorem was proven in [1]:

Theorem 3 ([1])

$$f_d(m) = g_d(m)$$

In the next section we give a natural one-to-one correspondence between polynomials and hypergraphs, proving Theorem 3.

For p prime, and α positive integer it is known ([1], [2], [4]) that $g_d(p^\alpha) = d(p^\alpha - 1) + 1$, so

Corollary 4 ([1]) *For $\mathcal{F} = \{F_1, F_2, \ldots, F_{d(q-1)+1}\}$, where q is a prime-power, and d denotes the maximal degree of the hypergraph, there exists an $\emptyset \neq \mathcal{F}_0 \subset \mathcal{F}$, such that $|\bigcup_{F \in \mathcal{F}_0} F| \equiv 0 \pmod{q}$.*

This corollary is a generalization of the undergraduate exercise that from arbitrary m integers, one can choose a non-empty subset, which adds up to 0 modulo m (the $d = 1$ case).

In 1991, *Barrington, Beigel* and *Rudich* [3] gave an explicit construction for polynomials modulo $m = p_1^{\alpha_1} p_2^{\alpha_2} \ldots p_r^{\alpha_r}$, showing that

$$g_d(m) = \Omega(d^r).$$

Since the proof of Theorem 3 (both the original and ours in the next section) gives explicit constructions for hypergraphs from polynomials, the following corollary holds:

Corollary 5 *Let $m = p_1^{\alpha_1} p_2^{\alpha_2} \ldots p_r^{\alpha_r}$. Then there exists an explicitly constructible hypergraph $\mathcal{F}$ of maximum degree d, such that $|\mathcal{F}| = \Omega(d^r)$ and for each $\emptyset \neq \mathcal{F}_0 \subset \mathcal{F}$ it is satisfied that $|\bigcup_{F \in \mathcal{F}_0} F| \not\equiv 0 \pmod{m}$.*

The authors of [1] gave a doubly-exponential upper bound to $f_d(m)$, which was based on a Ramsey-theoretic bound of [2]. More recently, *Tardos* and *Barrington* [4] showed, that

$$f_d(m) = \exp(O(d^{r-1})).$$

2 Correspondence between polynomials and hypergraphs

We give here a short and direct proof for Theorem 3. Let Q denote the set of rationals. It is well known, that the set of functions $\{f : \{0,1\}^t \to Q\}$ forms a 2^n-dimension vector space over the rationals. One useful basis of this vectorspace is the set of OR-functions $\{\bigvee_{i\in I} x_i : I \subset \{1,2,\ldots,n\}\}$, where

$$\bigvee_{i\in I} x_i = 1 - \prod_{i\in I}(1-x_i).$$

It is easy to see, that any integer-valued function on the hypercube can be written as the integer-coefficient linear combination of these OR-functions. Moreover, if the function is a degree-d polynomial, then it is enough to use OR functions with $|I| \leq d$. If we consider modulo m polynomials, then the coefficients can be restricted to the set $\{0,1,2,\ldots,m-1\}$. It will be convenient to view modulo m polynomials as the sum of several OR functions with coefficient 1; instead of multiplying an OR function with a coefficient a we will add it up exactly a times.

Consequently, our degree-d modulo m polynomial has the following form:

$$h = S_1 + S_2 + \cdots + S_\ell, \tag{1}$$

where S_i is an OR-function of degree at most d.

Now we are ready to define the one-to-one correspondence between degree-d modulo m polynomials without non-trivial zeroes on the hypercube and and hypergraphs, without non-empty sub-hypergraphs of modulo-m sum of 0. Let h be a degree-d polynomial written in form (1), and define hypergraph $\mathcal{F} = \{F_1, F_2, \ldots, F_t\}$, where $F_i = \{S_j : x_i \text{ appears as a variable in } S_j\}$. Clearly, the degree of this hypergraph is at most the degree of h, that is, d.

On the other hand, for a hypergraph $\mathcal{F} = \{F_1, F_2, \ldots, F_t\}$ on the ground-set $\{v_1, v_2, \ldots, v_\ell\}$, let us define $h(x_1, x_2, \ldots, x_t) = S_1 + S_2 + \cdots + S_\ell$, where

$$S_i = \bigvee_{j : v_i \in F_j} x_j.$$

Obviously, the degree of h is at most the degree of $\mathcal{F}$.

Clearly, $\mathcal{F}$ has a non-empty sub-hypergraph with union-size 0 modulo m if and only if there exists a $0 \neq x : h(x) \equiv 0 \pmod m$. To prove this, it is enought to see that $I = \{i : x_i = 1\}$ is the same I for which $|\bigcup_{i\in I} F_i| \equiv 0 \pmod m$.□

References

[1] N. Alon, D. Kleitman, R. Lipton, R. Meshulam, M. Rabin, and J. Spencer. Set systems with no union of cardinality 0 modulo m. *Graphs and Combinatorics*, 7:97–99, 1991.

[2] R. Baker and W. Schmidt. Diophantine problems in variables restricted to the values 0 and 1. *Journal of Number Theory*, 12:460–486, 1980.

[3] D. A. M. Barrington, R. Beigel, and S. Rudich. Representing Boolean functions as polynomials modulo composite numbers. *Comput. Complexity*, 4:367–382, 1994. Appeared also in *Proc. 24th Ann. ACM Symp. Theor. Comput.*, 1992.

[4] G. Tardos and D. A. M. Barrington. A lower bound on the MOD 6 degree of the OR function. *Comput. Complexity*, 7:99–108, 1998.